



NATIONAL MISSION FOR CLEAN GANGA [NMCG]

Department of Water Resources, River Development & Ganga Rejuvenation
Ministry of Jal Shakti, Government of India

VOLUME – I

INTEGRATED RIVER ASSESSMENT FRAMEWORK – SCIENTIFIC FOUNDATIONS, CLASSIFICATION AND ASSESSMENT METHODOLOGIES

Prepared By :



**Centre for Ganga River Basin
Management and Studies**
Indian Institute of Technology Kanpur



October 2026

© cGanga & NMCG, 2026

National Mission for Clean Ganga (NMCG)

NMCG is the implementation wing of National Ganga Council which was setup in October 2016 under the River Ganga Authority order 2016. Initially NMCG was registered as a society on 12th August 2011 under the Societies Registration Act 1860. It acted as implementation arm of National Ganga River Basin Authority (NGRBA) which was constituted under the provisions of the Environment (Protection) Act (EPA) 1986. NGRBA has since been dissolved with effect from the 7th October 2016, consequent to constitution of National Council for Rejuvenation, Protection and Management of River Ganga (referred to as National Ganga Council).

www.nmcg.in

Centre for Ganga River Basin Management and Studies (cGanga)

cGanga is a think tank formed under the aegis of NMCG, and one of its stated objectives is to make India a world leader in river and water science. The Centre is headquartered at IIT Kanpur and has representation from most leading science and technological institutes of the country. cGanga's mandate is to serve as think-tank in implementation and dynamic evolution of Ganga River Basin Management Plan (GRBMP) prepared by the Consortium of 7 IITs. In addition to this it is also responsible for introducing new technologies, innovations and solutions into India.

www.cganga.org

Acknowledgment

This document is a collective effort of many experts, scientific staff and students associated with cGanga, IIT Kanpur from multiple institutions. Contributions to the photographs and images by individuals are gratefully acknowledged.

Suggested Citation

cGanga & NMCG, 2026, Volume I: Integrated River Assessment Framework - Scientific Foundations, Classification and Assessment Methodologies

Contacts

Centre for Ganga River Basin Management and Studies (cGanga), Indian Institute of Technology Kanpur, Kanpur 208 016, Uttar Pradesh, India

Author

Vinod Tare

Former Professor, Founder & Advisor, cGanga, IIT Kanpur

VOLUME - I

**INTEGRATED RIVER ASSESSMENT FRAMEWORK -
SCIENTIFIC FOUNDATIONS, CLASSIFICATION AND
ASSESSMENT METHODOLOGIES**

October 2026



**Centre for Ganga River Basin Management and Studies
Indian Institute of Technology Kanpur**



Preface

Rivers are dynamic and interconnected ecological systems that sustain biodiversity, regulate hydrological processes, support livelihoods, and provide essential ecosystem services. India's diverse river systems, ranging from glacier-fed Himalayan rivers and monsoon-driven alluvial rivers to peninsular, arid, groundwater-fed and coastal systems, exhibit distinct natural characteristics and ecological functions. Their effective conservation and management therefore require a scientific approach that recognizes this diversity while distinguishing natural variability from human-induced degradation.

Conventional river assessment approaches have often emphasized water quality and selected environmental indicators, providing an incomplete understanding of the physical, ecological and functional processes that sustain river systems. Recognizing this limitation, the Integrated River Assessment Framework (IRAF) has been developed by the Centre for Ganga River Basin Management and Studies (cGanga), IIT Kanpur, within the broader context of the Ganga River Basin Management Plan (GRBMP 2.0). The framework advances an integrated, process-based approach to river assessment, conservation, restoration and sustainable management.

The present report brings together four complementary assessment components: the River Classification Framework (RCF), which establishes the intrinsic identity of river reaches; the Pristine River Grading Index (PRGI), which evaluates their intrinsic ecological significance; the River Health Assessment (RHA), which assesses their present functional condition relative to appropriate reference conditions; and the Functional Assessment and Classification of Local Water Systems (FACLWS), which evaluates the functions and functional deficits of associated streams, drains, ponds, wetlands, floodplain channels and other local water systems. Together, these components provide a comprehensive scientific basis for understanding rivers and their supporting aquatic networks.

A fundamental principle of IRAF is that river identity, ecological significance and present health are distinct but interrelated dimensions. By assessing each river reach in relation to its natural characteristics, the framework seeks to avoid misleading comparisons between fundamentally different river types and supports more appropriate, evidence-based management decisions.

The report further demonstrates how scientific assessment can be translated into practical action through Planning and Management of Local Water Systems (PMLWS) and reach-specific river-management responses. It establishes a structured basis for conservation prioritization, restoration planning, environmental flow assessment, floodplain management, urban river rejuvenation, ecological rehabilitation, and long-term monitoring.

Ultimately, IRAF represents a step towards a more holistic and scientifically grounded approach to river basin management - one that recognizes rivers as living systems and seeks to secure Water Security, River Health and Climate Resilience for present and future generations.

Dr Vinod Tare

Former Professor, Mentor & Advisor
Centre for Ganga River Basin Management & Studies (cGanga), IIT Kanpur



Content

	Page
Preface	5
Executive Summary	11
1 Introduction and Conceptual Foundation	16
1.1 Background	16
1.2 Need for a Unified River Assessment Framework	17
1.3 Philosophy of River Assessment	18
1.4 Rivers as Dynamic Functional Systems	19
1.5 Natural Variability and Anthropogenic Degradation	19
1.6 Integrated Assessment Framework	20
1.7 Scope and Organization of the Guideline	22
2 River Classification Framework	23
2.1 Introduction	23
2.2 Objectives of River Classification	23
2.3 Scientific Basis	24
2.4 Principles of River Classification	25
2.5 Components of River Classification	25
2.6 Climatic Setting	27
2.7 Longitudinal River Continuum	27
2.8 Integration with River Styles	28
2.9 Integrated River Classification Code	28
2.10 Applications of River Classification	29
2.11 Concluding Remarks	29
3 Pristine River Grading Index (PRGI)	30
3.1 Introduction	30
3.2 Purpose and Objectives	30
3.3 Conceptual Framework	31
3.4 Guiding Principles	31
3.5 Assessment Framework	32
3.6 PRGI Assessment Indicators	34
3.7 Weighting Scheme	34
3.8 Scoring Methodology	35
3.9 PRGI Grades	36
3.10 Interpretation of PRGI	36
3.11 Applications	37
3.12 Relationship with Other Assessment Frameworks	37
3.13 Limitations	37
3.14 Concluding Remarks	37
4 River Health Assessment (RHA)	39
4.1 Introduction	39
4.2 Purpose and Objectives	39
4.3 Conceptual Framework	40
4.4 Guiding Principles	40
4.5 Functional Dimensions of River Health	41
4.6 Integrated Functional Assessment Framework	44
4.7 Assessment Indicators	44
4.8 RHA Assessment Indicators and Weighting Scheme	44
4.9 Assessment Indicators	48
4.10 Scoring Methodology	49
4.11 River Health Classes	49

	Page
4.12 Interpretation of Health Classes	50
4.13 Management Decision Framework	51
4.14 Integration with River Classification and PRGI	51
4.15 Applications	52
4.16 Strengths and Limitations	52
4.17 Concluding Remarks	53
5 Functional Assessment and Classification of Local Water Systems (FACLWS)	54
5.1 Introduction	54
5.2 Need for Utilizing Local Water System to Develop Ecological Infrastructure	55
5.3 Objective	56
5.4 Guiding Principles	56
5.5 Local Water Systems	56
5.6 Functional Dimensions	58
5.7 Assessment Indicators	60
5.8 Functional Performance Categories	60
5.9 Management Strategies	60
5.10 Functional Assessment Matrix	61
5.11 Integration with RHA	61
5.12 Applications	62
5.13 Relationship with the Integrated Assessment Framework	62
5.14 Concluding Remarks	63
6 Integration into River Basin Management	64
6.1 Introduction	64
6.2 From Assessment to Management	64
6.3 Integrated River Assessment Hierarchy	65
6.4 Decision Matrix for River Basin Management	65
6.5 River Basin Zonation	66
6.6 Environmental Flow Management	66
6.7 River Restoration Planning	65
6.8 Nature-Based Solutions	67
6.9 Urban River Rejuvenation	68
6.10 Climate Change Adaptation	68
6.11 Basin Governance	68
6.12 Monitoring and Adaptive Management	69
6.13 Integrated River Basin Management Framework	69
6.14 Integration with GRBMP 2.0	70
6.15 Concluding Remarks	70
7 Longitudinal River Classification and Reach-Based Assessment	71
7.1 Introduction	71
7.2 Rationale for Reach-Based Assessment	71
7.3 Concept of River Reaches	72
7.4 Hierarchical Spatial Framework	72
7.5 Criteria for Reach Delineation	77
7.6 Reach Characterization	73
7.7 Integrated Reach Assessment	74
7.8 Longitudinal River Profile	74
7.9 Reach Prioritization	75
7.10 Reach Report Card	75
7.11 GIS-Based River Assessment	75
7.12 Applications	76
7.13 Integration with GRBMP 2.0	76

7.14	Concluding Remarks	76
8	Assessment Methodology and Implementation Protocols	77
8.1	Introduction	77
8.2	Implementation Framework	76
8.3	Stage I – Preliminary Desk Study	78
8.4	Stage II – River Segmentation	78
8.5	Stage III – Field Investigation	78
8.6	Stage IV – Data Compilation and Validation	79
8.7	Stage V – Assessment Procedures	80
8.8	Scoring Protocol	80
8.9	Data Quality Assurance	81
8.10	GIS-Based Assessment System	81
8.11	River Reach Report Card	82
8.12	Basin Summary Dashboard	82
8.13	Monitoring Frequency	82
8.14	Capacity Building	83
8.15	Reporting Standards	83
8.16	Digital Decision Support System	84
8.17	Concluding Remarks	84
9	Implementation Framework for Integrated River Basin Management	85
9.1	Introduction	85
9.2	From Assessment to Integrated River Basin Management	88
9.3	Guiding Principles for Implementation	90

Concluding Note



INTEGRATED RIVER ASSESSMENT FRAMEWORK - SCIENTIFIC FOUNDATIONS, CLASSIFICATION AND ASSESSMENT METHODOLOGIES

Executive Summary

Background

India possesses one of the world's most diverse river systems, extending from glacier-fed Himalayan rivers to monsoon-driven peninsular rivers, groundwater-fed spring systems, ephemeral desert streams, and tidally influenced estuaries. These rivers differ fundamentally in their hydrology, geomorphology, sediment dynamics, ecology, hydrochemistry, and ecosystem functions. Consequently, no single indicator or uniform assessment standard can adequately represent the condition, significance, or management requirements of all rivers. Conventional river assessment methods have largely focused on water quality, often treating rivers as conduits of water rather than dynamic ecological systems. Such approaches inadequately capture the complex interactions among hydrological, geomorphological, ecological, and biogeochemical processes that sustain river health. They also fail to distinguish natural variability from anthropogenic degradation, leading to inappropriate comparisons among fundamentally different river types.

Recognizing these limitations, cGanga, IIT Kanpur, under the broader framework of the **Ganga River Basin Management Plan (GRBMP 2.0)**, has developed a unified and scientifically robust framework that integrates **River Classification (RCF)**, **Pristine River Grading Index (PRGI)**, **River Health Assessment (RHA)**, and **Functional Assessment and Classification of Local Water Systems (FACLWS)**. The outputs of RCF, PRGI, RHA and FACLWS are synthesised through Integrated Diagnosis and translated into two complementary management pathways: Reach-Specific River Management and Planning and Management of Local Water Systems (PMLWS). Implementation is followed by monitoring, periodic reassessment and adaptive management.

Conceptual Foundation

The framework is founded on three fundamental principles.

First, every river, more precisely every river reach, possesses intrinsic ecological value irrespective of its size, water clarity, or economic importance. River stretches differ naturally because of their geology, climate, hydrology, sediment regime, and evolutionary history. These natural differences must be recognized rather than interpreted as degradation.

Second, river assessment should distinguish between intrinsic characteristics and current condition. A naturally sediment-rich Himalayan River and a clear groundwater-fed spring

stream are equally natural systems, although they differ greatly in appearance and water chemistry.

Third, river basin management must adopt a systems perspective in which rivers, floodplains, wetlands, streams, ponds, drains, and groundwater-dependent ecosystems function as interconnected components of a single hydrological and ecological network.

Accordingly, the framework moves beyond conventional water quality assessment towards an integrated understanding of rivers as dynamic, living systems.

Four-Component Integrated Assessment Framework

The Integrated River Assessment Framework comprises four complementary assessment and diagnostic components: RCF, PRGI, RHA and FACLWS. Their outputs are synthesised through Integrated Diagnosis, which provides the scientific basis for Reach-Specific River Management and PMLWS. Neither management pathway constitutes a fifth assessment framework.

I. River Classification Framework: Question addressed → What type of river is it?

The first step is to identify the intrinsic characteristics of a river through a multidimensional classification system. Classification is independent of river condition and represents the natural identity of the river.

The framework classifies rivers using five principal descriptors:

- Hydrological regime
- Geomorphological setting
- Sediment regime
- Hydrochemical characteristics
- Source of flow

Additional descriptors include climatic setting, longitudinal position, and river style.

Each river reach is assigned a standardized classification code that provides a concise representation of its intrinsic characteristics, thereby facilitating consistent documentation, comparison, and basin-wide analysis.

II. Pristine River Grading Index (PRGI): Question addressed → How ecologically important is the river under natural conditions?

The Pristine River Grading Index evaluates the intrinsic ecological significance of rivers or river reaches without considering anthropogenic disturbance.

The assessment is based on four pillars:

- Hydrology
- Biodiversity
- Uniqueness

- Ecosystem Services

Each pillar is quantified through measurable indicators and integrated into a weighted scoring framework.

The resulting PRGI Tiers identify rivers, or river reaches of exceptional ecological significance, thereby supporting conservation prioritization, protected area planning, biodiversity management, and ecological safeguarding.

Importantly, PRGI evaluates ecological importance rather than environmental quality. Naturally turbid, sediment-rich, or mineralized rivers are therefore assessed according to their natural characteristics and are not penalized for conditions that arise from natural processes.

III. **River Health Assessment (RHA): Question addressed → What is the present river-health condition of the river or reach relative to its appropriate reference condition?**

RHA evaluates six integrated components: Hydrological Condition (H), Geomorphic Condition (G), Sediment Regime (S), Water & Environmental Quality (W), Habitat & Biodiversity (B), and Connectivity (C).

Unlike assessments based solely on water quality, RHA evaluates the present condition of a river or identified reach through six interrelated hydrological, geomorphic, sediment, water-quality, biodiversity and connectivity components.

Assessment is undertaken relative to an appropriate reference condition for each river type, thereby distinguishing natural variability from anthropogenic alteration. River reaches are assigned one of the five River Health Classes:

A – Pristine/Near-reference; B – Slightly Modified; C – Moderately Altered; D – Heavily Altered; E – Critically Degraded.

RHA provides a scientifically defensible basis for restoration prioritization, environmental flow assessment, ecological rehabilitation, and basin planning.

IV. **Functional Assessment and Classification of Local Water Systems (FACLWS): What functions should associated local water systems perform, and how well are they functioning?**

The framework extends beyond main river channels to include streams, drains, ponds, wetlands, floodplain channels, and other local water bodies.

Recognizing that many urban and peri-urban systems can no longer realistically be restored to pristine conditions, the framework evaluates these systems according to their functional performance rather than their degree of naturalness.

Assessment considers five functional dimensions:

- Hydrological functions
- Water-quality functions
- Ecological functions
- Geomorphic/sediment functions
- Human-support functions

Habitat and ecological connectivity are assessed within Ecological functions, biogeochemical processes within Water-quality functions and compatible uses, livelihoods socio-cultural benefits within Human-support functions.

Delete the first occurrence of this paragraph. Retain its second occurrence under the heading Integration within River Basin Management.

Integration within River Basin Management

River Classification establishes river/reach identity and reference context. PRGI establishes intrinsic ecological significance. RHA diagnoses present river-health condition. FACLWS diagnoses the functions and functional deficits of associated Local Water Systems. These diagnostic outputs are integrated and translated into action through PMLWS and reach-specific river-management responses.

Together, these assessments enable managers to distinguish between rivers requiring protection, restoration, rehabilitation, or sustainable management.

Applications

The framework supports a broad range of river basin management activities, including:

- Conservation prioritization
- River restoration planning
- Environmental flow assessment
- Basin zoning and land-use planning
- Floodplain management
- Urban river rejuvenation
- Nature-based wastewater treatment
- Biodiversity conservation
- Climate resilience planning
- Long-term river health monitoring

The framework is equally applicable to Himalayan, alluvial, peninsular, coastal, arid, and groundwater-fed river systems across India.

Conclusion

Rivers are dynamic, interconnected ecological systems whose health depends upon the integrity of hydrological, geomorphological, ecological, and biogeochemical processes operating across entire river basins. Sustainable river management therefore requires assessment frameworks that recognize both natural diversity and anthropogenic impacts.

The unified framework presented provides a scientifically robust and operationally practical basis for understanding river identity, ecological significance, functional health, and basin-wide ecological performance. By integrating river classification, ecological grading, functional health assessment, and functional evaluation of local water systems, the framework establishes a comprehensive foundation for conservation, restoration, and sustainable management of India's river systems.

The intended integrated outcomes are Water Security, River Health and Climate Resilience.

1. Introduction and Conceptual Foundation

1.1. Background

Rivers are among the most dynamic and complex natural systems on Earth. They are not merely channels conveying water from upstream to downstream, but integrated ecological systems in which hydrological, geomorphological, geological, chemical, and biological processes continuously interact across space and time. These interactions shape river morphology, regulate ecosystem functions, sustain biodiversity, recharge groundwater and receive base flows through subsurface flows, transport sediments and nutrients, and provide a wide range of ecosystem services essential for environmental sustainability and human well-being.

India possesses extraordinary river diversity. The country encompasses glacier-fed Himalayan rivers, snow-fed mountain streams, monsoon-driven alluvial rivers, groundwater-fed spring systems, peninsular plateau rivers, ephemeral desert streams, tidal rivers, estuaries, and distributary networks. Each river type has evolved under distinct geological, climatic, and hydrological settings and therefore exhibits unique physical, chemical, and ecological characteristics.

Historically, river management has largely been guided by sectoral objectives such as irrigation, flood control, navigation, hydropower development, urban water supply, wastewater disposal, and pollution control. While these interventions have supported socio-economic development, they have often overlooked the ecological processes that sustain river systems. As a result, many rivers have experienced altered flow regimes, disrupted sediment transport, floodplain disconnection, habitat degradation, declining biodiversity, and deteriorating water quality.

In recent decades, increasing recognition of rivers as living ecosystems has fundamentally changed the philosophy of river management. Contemporary approaches emphasize ecological integrity, process-based understanding, environmental flows, nature-based solutions, and integrated river basin management. Rather than focusing solely on water quality or infrastructure, these approaches seek to maintain or restore the natural functions that enable rivers to remain resilient and self-sustaining.

The **Ganga River Basin Management Plan (GRBMP 2.0)** framework developed by cGanga, IIT Kanpur adopts this broader perspective. It recognizes that sustainable river management requires an integrated framework capable of distinguishing a river's intrinsic natural characteristics from its present condition, while also acknowledging the contribution of associated water systems such as streams, wetlands, ponds, floodplain channels, and urban drains as local ecological infrastructure.

1.2. Need for a Unified River Assessment Framework

Most existing river assessment methods evaluate only selected aspects of river systems. Water quality indices focus primarily on physico-chemical parameters; biological indices emphasize aquatic communities; hydrological assessments examine flow regimes; geomorphological studies investigate channel forms and sediment processes; and ecological assessments often address habitat or biodiversity independently. Although each provides valuable information, none individually represents the overall condition or functioning of an entire river system.

An additional limitation arises from the widespread use of uniform assessment criteria across fundamentally different river types. Glacier-fed rivers naturally exhibit high sediment loads and seasonal variability, while spring-fed rivers generally have clearer water and more stable discharge. Likewise, alluvial rivers differ significantly from bedrock-confined mountain rivers in morphology, habitat diversity, and sediment transport. Evaluating such diverse systems using identical standards can lead to misleading conclusions and inappropriate management decisions.

Another common limitation is the failure to distinguish **intrinsic ecological significance** from **current condition**. A river reach of exceptional ecological importance may presently be degraded, while another reach with relatively modest ecological significance may remain in excellent condition. These represent two different management questions requiring different assessment approaches.

Furthermore, local water systems—including urban drains, natural streams, ponds, wetlands, floodplain channels, and decentralized treatment systems—are often assessed independently of the river basin despite their important ecological and hydrological linkages with the main river channel.

These limitations underscore the need for a unified assessment framework that:

- Recognizes natural diversity among river systems.
- Separates river identity from river condition.
- Integrates ecological, hydrological, geomorphological, and functional perspectives.
- Supports consistent river basin planning.
- Provides practical guidance for conservation, restoration, and sustainable management.

The framework presented in this guideline addresses these requirements through four complementary assessment components:

River Classification Framework (RCF), Pristine River Grading Index (PRGI), River Health Assessment (RHA), and Functional Assessment and Classification of Local Water Systems (FACLWS).

1.3. Philosophy of River Assessment

The framework is founded on the principle that **every river should be evaluated in relation to its own natural potential rather than against a universal standard**. Rivers are products of long-term interactions among geology, climate, hydrology, topography, sediment dynamics, and ecological evolution. Their natural characteristics therefore differ considerably across regions and river types.

A scientifically meaningful assessment must first establish what a river is naturally before evaluating how its present condition has changed. This distinction between **natural character** and **current condition** is fundamental to the framework.

Accordingly, the assessment philosophy recognizes four complementary assessment questions:

1. What type of river is it?

This is addressed through River Classification, which identifies the river's intrinsic bio-physical and hydrological characteristics.

2. How important is the river under natural conditions?

This is evaluated using the Pristine River Grading Index (PRGI), which measures intrinsic ecological significance and conservation value.

3. How well is the river functioning currently relative to its natural reference condition?

This is assessed through the River Health Assessment (RHA), which evaluates the integrity of hydrological, geomorphological, ecological, hydrochemical, and connectivity processes.

4. What functions should associated Local Water Systems perform, how well are they presently functioning, and how do their functional deficits influence the receiving river? FACLWS addresses this question by diagnosing the expected functions, present functional condition, deficits and river influence of individual Local Water Systems and their connected networks?

This is addressed through the potential of Local Water Systems to be a part of the ecological infrastructure in sync with the natural processes and systems, recognizing the contribution of streams, wetlands, ponds, drains, and floodplain channels to basin-wide resilience.

These four questions constitute assessment and diagnosis. Their outputs are synthesised through Integrated Diagnosis to identify management requirements for two coordinated pathways: Reach-Specific River Management and PMLWS.

1.4. Rivers as Dynamic Functional Systems

A river is best understood as a **dynamic functional continuum** rather than a static watercourse. From its source to its outlet, a river continuously evolves in response to changes in climate, topography, geology, sediment supply, tributary inflows, groundwater interactions, and human interventions.

Hydrological processes regulate the quantity, timing, duration, and variability of water flows. Geomorphological processes shape channel morphology through erosion, transport, and deposition of sediments. Hydrochemical processes govern nutrient cycling, dissolved constituents, and water quality. Ecological processes sustain biodiversity, food webs, habitat complexity, and ecosystem resilience. Connectivity processes link upstream and downstream reaches, rivers with floodplains, surface water with groundwater, and aquatic ecosystems with surrounding landscapes.

These processes are highly interdependent. Alterations in one component frequently propagate through the entire river system. For example, interruption of sediment transport by dams may trigger downstream channel incision, habitat simplification, floodplain disconnection, and ecological degradation. Similarly, excessive abstraction of river flows can affect groundwater recharge, wetland hydrology, water quality, and biodiversity.

Consequently, meaningful river assessment requires evaluation of the functioning of the river as an integrated socio-ecological system rather than isolated measurements of individual parameters.

1.5. Natural Variability and Anthropogenic Degradation

One of the most important principles underlying this guideline is the distinction between **natural variability** and **human-induced degradation**.

Natural variability reflects the inherent diversity among rivers arising from differences in climate, geology, geomorphology, hydrology, vegetation, and landscape evolution. Such variability should not be interpreted as environmental degradation. For example:

- Glacier-fed Himalayan rivers naturally carry high sediment loads and exhibit seasonal fluctuations associated with snowmelt and monsoon rainfall.
- Peninsular rivers generally experience stronger seasonal flow variability linked to monsoon rainfall.
- Spring-fed rivers often maintain relatively stable discharge throughout the year.
- Arid-zone rivers may remain dry for extended periods as a natural characteristic rather than an indication of ecological failure.
- Estuarine rivers naturally experience tidal influence and elevated salinity.

These characteristics represent natural river behaviour and therefore define appropriate reference conditions for assessment.

Anthropogenic degradation, in contrast, results from alterations such as flow regulation, excessive water abstraction, pollution, floodplain encroachment, channel modification, sediment trapping, riparian vegetation loss, habitat fragmentation, invasive species, and unsustainable land-use practices.

The framework therefore evaluates rivers relative to the conditions expected for their specific river type rather than against generalized standards.

1.6. Integrated Assessment Framework

The unified framework consists of four complementary assessment and diagnostic components. River Classification Framework (RCF) establishes river/reach identity and reference context using Hydrology, Geomorphology, Sediment, Hydrochemistry and Source; climatic setting, longitudinal position and River Styles are supplementary descriptors, and anthropogenic regulation is recorded separately as a modifier. PRGI assesses intrinsic ecological significance. RHA assesses present river-health condition through H, G, S, W, B and C relative to the appropriate reference condition. FACLWS assesses the expected functions, present functional condition, deficits and river influence of associated Local Water Systems. Integrated Diagnosis synthesises these outputs. Reach-Specific River Management and PMLWS are distinct, coordinated planning and management pathways, followed by implementation, monitoring, reassessment and adaptive management.

- **River Classification**

Defines the intrinsic identity of the river based on hydrology, geomorphology, sediment regime, hydrochemistry, source of flow, and longitudinal characteristics.

- **Pristine River Grading Index (PRGI)**

Evaluates the intrinsic ecological significance of rivers under natural conditions, independent of present degradation.

- **River Health Assessment (RHA)**

Assesses the current functional condition of rivers relative to their natural reference state by integrating hydrological, geomorphological, hydrochemical, ecological, habitat, connectivity, sediment, and ecosystem service dimensions.

- **Local Water Systems**

The associated water systems—including streams, drains, ponds, wetlands, and floodplain channels—based on their ecological, hydrological, treatment, habitat, and socio-cultural functions serve as local ecological infrastructure.

Planning and Management of Local Water Systems (PMLWS):

Translates FACLWS diagnosis, together with relevant RHA requirements and planning information, into protection, sustenance, restoration, rehabilitation, reconnection, optimisation/repurposing and, where justified, augmentation of Local Water Systems.

Together, these four components and PMLWS form an integrated decision-support framework that enables:

- scientifically sound river assessment;
- conservation prioritization;
- restoration planning;
- environmental flow assessment;
- basin zoning and land-use planning;
- nature-based wastewater management;
- monitoring of river health and resilience; and
- adaptive river basin governance.

The overall conceptual framework adopted in this guideline is illustrated in Figure 1.1, which integrates the four complementary assessment components into a unified river basin management framework.




Figure 1.1: Integrated River Assessment Framework (IRAF) - From River Classification and Integrated Diagnosis to Planning, Management and Outcomes

River Classification (RCF) establishes river/reach identity and reference context; PRGI determines intrinsic ecological significance; RHA evaluates present river-health condition; and FACLWS diagnoses the functions and deficits of associated Local Water Systems. Integrated Diagnosis synthesises these outputs and informs two coordinated pathways: Reach-Specific

River Management and PMLWS. Implementation and monitoring provide evidence for reassessment and adaptive management, contributing to Water Security, River Health and Climate Resilience.

1.7. Scope and Organization of the Guideline

This guideline provides a comprehensive methodology for classifying, grading, assessing, and managing river systems within the context of integrated river basin management. While developed under the framework of the Ganga River Basin Management Plan (GRBMP 2.0), the concepts and methodologies are applicable to diverse river systems across India.

The chapters that follow present each component of the framework in a logical sequence. Chapter 2 introduces the River Classification Framework, establishing the natural identity and typology of river systems. Chapter 3 presents the Pristine River Grading Index (PRGI), followed by Chapter 4 on the River Health Assessment (RHA). Chapter 5 addresses the use and development of Local Water Systems as supporting ecological infrastructure, while subsequent chapters demonstrate how these frameworks are integrated into river basin planning, management, and decision-making.

By progressing from river identity to ecological significance, functional health, and basin-wide integration, the guideline offers a coherent and scientifically rigorous foundation for the conservation, restoration, and sustainable management of India's river systems.

2. River Classification Framework

2.1. Introduction

River classification forms the scientific foundation of integrated river basin management. Before a river can be evaluated for its ecological significance, functional health, or management priority, its intrinsic natural characteristics must first be understood. Rivers differ fundamentally in their hydrological regimes, geomorphological settings, sediment dynamics, hydrochemical properties, and sources of flow. These characteristics determine how rivers function, evolve, and respond to both natural processes and anthropogenic disturbances.

Historically, many river assessment frameworks have implicitly assumed that all rivers can be evaluated against common standards. Such approaches overlook the natural diversity of river systems and often lead to inappropriate comparisons. A glacier-fed Himalayan River carrying high suspended sediment loads, for example, should not be assessed using the same criteria as a spring-fed river of the Western Ghats or an ephemeral stream of western Rajasthan. Their contrasting characteristics reflect natural environmental settings rather than differences in ecological quality.

River classification therefore serves as the essential first step in river assessment. It establishes the natural identity of a river and provides the contextual basis for all subsequent evaluations. Within the framework presented in this guideline, river classification is independent of ecological condition or anthropogenic disturbance. It describes **what a river is**, not **how healthy it is**.

The classification framework presented here integrates hydrological, geomorphological, sedimentological, hydrochemical, climatic, and ecological perspectives into a standardized typology that is applicable across the diverse river systems of India.

2.2. Objectives of River Classification

The River Classification Framework has six principal objectives:

- To establish the intrinsic natural identity of river reaches.
- To distinguish natural river characteristics from anthropogenic alterations.
- To provide a standardized basis for ecological grading and functional health assessment.
- To facilitate comparison among rivers of similar types rather than dissimilar systems.
- To support river basin planning, conservation, restoration, and environmental flow assessment.
- To enable consistent documentation and geospatial mapping of river characteristics.

Rather than assigning quality grades, the framework provides a descriptive classification that captures the natural variability of Indian river systems. The overall hierarchy of the River Classification Framework is presented in Figure 2.1.




Figure 2.1: River Classification Framework

2.3. Scientific Basis

River systems evolve through the interaction of climate, geology, topography, hydrology, vegetation, and tectonic processes operating over geological timescales. These factors collectively determine river morphology, channel behaviour, sediment transport, hydrochemistry, ecological communities, and longitudinal evolution.

The classification framework adopts a **process-based approach**, recognizing that river form and function are inseparable. Hydrological processes govern discharge and flow variability; geomorphological processes shape channel morphology; sediment dynamics influence channel stability and habitat diversity; hydrochemical conditions regulate nutrient cycling and water chemistry; while ecological processes determine biodiversity and ecosystem functioning.

Accordingly, the framework integrates five primary descriptors:

- Hydrological Regime
- Geomorphological Setting
- Sediment Regime
- Hydrochemical Characteristics
- Source of Flow

These descriptors are supplemented by climatic setting, longitudinal river position, and River Styles to provide a comprehensive characterization of river systems.

2.4. Principles of River Classification

The classification framework is guided by the following principles:

Naturalness: Classification describes the river's natural characteristics rather than its present environmental condition.

Process-Based Understanding: Classification is based on dominant physical and ecological processes instead of isolated parameters.

Reach-Based Assessment: Classification is undertaken at the river reach level because river characteristics evolve continuously from source to mouth.

Hierarchical Organization: Broad River categories are progressively subdivided into increasingly specific river types.

Scalability: The framework can be applied to headwater streams, tributaries, major rivers, river basins, and national river inventories.

Compatibility: The classification supports integration with GIS databases, River Styles analysis, environmental flow studies, and ecological assessment frameworks.

2.5. Components of River Classification

The proposed framework classifies every river reach using five principal descriptors.

2.5.1. Hydrological Regime

Hydrology defines the quantity, timing, duration, frequency, and variability of river discharge. It controls channel-forming processes, habitat diversity, sediment transport, ecological connectivity, and biogeochemical cycling.

Table 2.1: River Hydrological Classes

Code	River Type	Characteristics
P	Perennial	Flow throughout the year
I	Intermittent	Seasonal cessation of flow
E	Ephemeral	Flow only after rainfall
T	Tidal	Flow influenced by tides

Where multiple conditions occur, the dominant natural regime should be recorded, with regulation indicated as a modifying attribute rather than a separate intrinsic hydrological class.

2.5.2. Geomorphological Setting

Geomorphology reflects the physical template within which river processes operate. It influences channel stability, floodplain development, sediment transport, habitat diversity, and ecological functioning.

Table 2.2: River Geomorphological Classes

Code	Setting
B	Bedrock
C	Colluvial
A	Alluvial
M	Mixed Bedrock–Alluvial
D	Deltaic
E	Estuarine

Within each setting, channel confinement may be described as:

- Confined
- Partly confined
- Laterally unconfined

This terminology aligns with the River Styles framework and supports process-based interpretation.

2.5.3. Sediment Regime

Sediment transport is a fundamental driver of river morphology and ecological processes. The sediment regime influences channel pattern, habitat complexity, floodplain evolution, and nutrient dynamics.

Table 2.3: Sediment Classes

Code	Sediment Condition
L	Low
M	Moderate
H	High
VH	Very High

Where possible, sediment characterization should incorporate both suspended and bed-load transport. Importantly, high sediment loads in Himalayan or braided alluvial rivers should be recognized as natural characteristics and not interpreted as ecological degradation.

2.5.4. Hydro-chemical Characteristics

Hydrochemistry reflects the natural interaction between water, geology, soils, climate, and groundwater. It influences aquatic ecology, nutrient availability, and suitability for various ecosystem functions.

Table 2.4: River Hydro-chemical Classes (Based on Total Dissolved Solids)

Code	TDS Level
L	Low
M	Moderate
H	High

Additional descriptors may include dominant ionic composition, alkalinity, hardness, pH, or salinity where appropriate. Natural hydrochemical characteristics should always be distinguished from pollution-related changes.

2.5.5. Source of Flow

The origin of river discharge strongly influences seasonal variability, hydrochemistry, sediment supply, and ecological dynamics.

Table 2.5: River Source Categories

Code	Source
G	Glacier-fed
S	Snow-fed
R	Rain-fed
W	Groundwater-fed
T	Tidal
C	Composite (mixed sources)

Many large rivers exhibit changing source dominance along their longitudinal profile. Accordingly, source classification should be applied reach-wise rather than assuming a single category for the entire river.

2.6. Climatic Setting

Climate exerts a dominant influence on river hydrology, sediment production, evapotranspiration, vegetation, and ecological productivity.

For management purposes, rivers may additionally be categorized according to their prevailing climatic setting:

- Himalayan alpine
- Humid subtropical
- Tropical monsoon
- Semi-arid
- Arid
- Coastal humid

This climatic descriptor provides valuable context for interpreting river behaviour and selecting appropriate reference conditions.

2.7. Longitudinal River Continuum

River characteristics evolve systematically from source to mouth, reflecting downstream changes in slope, discharge, sediment transport, floodplain development, hydrochemistry, and ecological communities.

The framework recognizes five generalized longitudinal zones:

Table 2.6: River Longitudinal Zonation

Zone	Dominant Characteristics
Headwaters	Steep gradients, confined channels, coarse sediment
Upper Reaches	Transition from erosion to transport
Middle Reaches	Active floodplains, sediment transport, diverse habitats
Lower Reaches	Deposition, meandering, distributaries
Estuarine/Tidal Zone	Tidal interaction, salinity gradients, delta formation

These zones should be interpreted flexibly, recognizing regional variability.

2.8. Integration with River Styles

The River Classification Framework is fully compatible with the **River Styles** approach developed by cGanga. River Styles provides a geomorphological understanding of river character based on valley setting, channel planform, confinement, bed material, and geomorphic units.

Within the present framework:

- River Styles describes the geomorphic character and behaviour of a river reach.
- River Classification provides the broader hydrological, sedimentological, hydro-chemical, and source context.
- Together, they establish the physical template upon which ecological grading and functional health assessments are undertaken.

The integration of River Styles strengthens process-based interpretation, supports prediction of river behaviour, and improves the design of conservation and restoration strategies.

2.9. Integrated River Classification Code

To facilitate standardized documentation, each river reach is assigned a concise alphanumeric code representing its dominant natural characteristics.

The recommended coding format is:

[Hydrology – Geomorphology – Sediment – Hydrochemistry – Source]

Table 2.7: Example River Reach Classification Codes

River Reach	Code	Interpretation
Glacier-fed Himalayan headwater	P–B–L–L–G	Perennial, bedrock-controlled, low sediment, low TDS, glacier-fed
Himalayan foothill reach	P–M–M–L–S	Perennial, mixed bedrock–alluvial, moderate sediment, low TDS, snow-fed
Alluvial plains	P–A–H–M–R	Perennial, alluvial, high sediment, moderate TDS, rain-fed
Groundwater-dominated reach	P–A–M–M–W	Perennial, alluvial, moderate sediment, moderate TDS, groundwater-fed
Tidal estuary	T–E–M–H–T	Tidal, estuarine, moderate sediment, high TDS, tidal influence

Anthropogenic regulation shall be recorded separately as a modifier and shall not replace intrinsic river identity. The coding system provides a compact yet comprehensive description that is readily incorporated into GIS databases, basin inventories, and river information systems.

2.10. Applications of River Classification

River classification supports a wide range of scientific and management applications, including:

- Establishing reference conditions for ecological assessment.
- Delineating river reaches for basin planning.
- Supporting the Pristine River Grading Index (PRGI).
- Providing the baseline for the River Health Assessment (RHA).
- Identifying environmentally sensitive river types.
- Guiding environmental flow assessments.
- Supporting River Styles analyses.
- Developing national river inventories.
- Standardizing communication among scientists, planners, and policymakers.

2.11. Concluding Remarks

River classification represents the essential first step in understanding and managing river systems.

By defining the intrinsic identity of river reaches, it establishes the scientific foundation upon which ecological grading, functional health assessment, and basin planning are built.

The integrated framework presented in this chapter recognizes the diversity of Indian river systems while providing a standardized and nationally applicable classification methodology. By combining hydrology, geomorphology, sediment dynamics, hydrochemistry, source of flow, climatic context, and River Styles, it offers a robust and process-based characterization of rivers that is directly applicable to conservation, restoration, and sustainable river basin management.

3. Pristine River Grading Index (PRGI)

3.1. Introduction

River conservation begins with recognizing the **intrinsic ecological significance** of rivers before evaluating their present condition. While many assessment frameworks focus on degradation, pollution, or restoration needs, equally important is the identification of rivers and river reaches that possess exceptional natural value and therefore warrant the highest levels of protection.

Every river has evolved under unique geological, hydrological, climatic, and ecological conditions. Some rivers support rare habitats, endemic species, intact floodplains, or unique geomorphic features, while others play critical roles in groundwater recharge, sediment transport, nutrient cycling, climate regulation, or cultural heritage. These characteristics collectively define the **ecological importance** of a river independent of its current environmental condition.

The **Pristine River Grading Index (PRGI)** has been developed to quantify this intrinsic ecological significance. Unlike pollution indices or river health assessments, PRGI does **not** evaluate degradation. Instead, it identifies the ecological value that would exist under natural or minimally disturbed conditions. Consequently, a highly significant river may presently be degraded, while a river of relatively lower ecological significance may currently remain in excellent condition.

PRGI therefore provides the scientific basis for conservation prioritization and serves as the second tier of the integrated river assessment framework presented in this guideline.

3.2. Purpose and Objectives

The primary purpose of PRGI is to evaluate the natural ecological significance of river systems. The specific objectives are:

- To identify rivers and rivers' reaches possessing exceptional ecological importance.
- To provide an objective basis for conservation prioritization.
- To distinguish intrinsic ecological value from present environmental condition.
- To support basin planning and protected area management.
- To identify ecologically sensitive river reaches requiring long-term protection.
- To provide baseline information for sustainable development planning.
- To complement the River Health Assessment (RHA).

PRGI is therefore a **conservation assessment tool**, not a pollution or river health index.

3.3. Conceptual Framework

The conceptual basis of PRGI is founded on a simple but fundamental principle: **Ecological significance is an inherent property of a river system and must be assessed independently of its present level of degradation.**

This distinction is essential because management objectives differ depending on whether the assessment concerns:

- intrinsic ecological importance, or
- present functional condition.

For example:

A glacier-fed Himalayan River supporting unique alpine biodiversity may possess extremely high ecological significance even if downstream reaches have experienced anthropogenic alteration.

A river may improve its RHA condition following restoration without changing its PRGI Tier. Urban setting, present modification and water-quality improvement do not, by themselves, establish intrinsic ecological significance; PRGI is determined from the river's natural hydrology, biodiversity, uniqueness and ecosystem services.

PRGI therefore answers the question:

"How important is this river in its natural state?"

rather than

"How healthy is this river at any instance?"

The latter question is addressed separately through the River Health Assessment (RHA).

3.4. Guiding Principles

The PRGI framework is governed by the following principles.

Natural Reference: Assessment is based on natural or least-disturbed ecological conditions rather than current anthropogenic influences.

Ecological Integrity: The index evaluates the ability of river systems to sustain naturally functioning ecosystems and ecological processes.

Multi-dimensional Assessment: Ecological significance cannot be represented by a single variable. Multiple ecological dimensions are therefore evaluated simultaneously.

Scientific Objectivity: Indicators are measurable, reproducible, and based on accepted scientific understanding.

River-Type Specificity: Assessment is undertaken relative to the natural characteristics of each river type.

Reach-Based Assessment: Large rivers should be assessed at the reach level because ecological significance changes along the river continuum.

3.5. Assessment Framework

The PRGI evaluates four fundamental dimensions of ecological significance.

3.5.1. Hydrological Importance

Hydrology provides the foundation upon which river ecosystems develop. Hydrological importance reflects the natural flow characteristics that sustain ecological processes and basin functioning.

Representative indicators include:

- natural flow regime
- perenniality
- environmental flow reliability
- groundwater interaction
- floodplain connectivity
- hydrological variability
- natural recharge functions

Hydrological importance reflects the river's capacity to sustain long-term ecological processes.

3.5.2. Biodiversity Value

Biodiversity represents one of the most important indicators of ecological significance.

Assessment considers:

- species richness
- endemic species
- threatened species
- migratory species
- habitat diversity
- aquatic biodiversity
- riparian biodiversity
- ecological communities

The emphasis is placed on naturally occurring biodiversity rather than artificially introduced species.

3.5.3. Uniqueness

Some rivers possess exceptional scientific or ecological value because of their rarity or distinctive natural characteristics.

Examples include:

- glacier-fed rivers
- spring-fed systems
- braided rivers
- karst rivers
- endemic fish habitats
- unique geomorphological features
- culturally significant natural landscapes

Uniqueness reflects irreplaceability rather than ecological condition.

3.5.4. Ecosystem Services

Healthy rivers provide numerous ecosystem services supporting both nature and society.

PRGI evaluates natural ecosystem services such as:

Regulating Services

- groundwater recharge
- flood moderation
- sediment regulation
- nutrient cycling
- climate regulation

Supporting Services

- habitat provision
- primary productivity
- ecological connectivity
- food web support

Provisioning Services

- freshwater availability
- fisheries
- natural resources

Cultural Services

- recreation
- education
- spiritual values
- cultural heritage
- aesthetic importance

Assessment emphasizes services provided under natural ecological functioning. The scheme followed for assigning river grades is illustrated in Figure 3.1.




Figure 3.1: Pristine River Grading Pyramid

3.6. PRGI Assessment Indicators

Each pillar is represented by a set of measurable indicators.

Table 3.1: PRGI Assessment Framework

Pillar	Representative Indicators	Recommended Weightages, %
Hydrology	Natural flow regime, perenniality, recharge, floodplain interaction	30
Biodiversity	Species richness, endemism, threatened species, habitat diversity	30
Uniqueness	Rare habitats, geomorphic uniqueness, endemic ecosystems	20
Ecosystem Services	Regulating, supporting, provisioning and cultural services	20

The indicator list may be refined according to basin-specific requirements while maintaining consistency with the conceptual framework.

3.7. Weighting Scheme

The four pillars contribute differently to intrinsic ecological significance. The recommended weighting is shown in Table 3.1. Hydrology and biodiversity receive the highest weight because they fundamentally determine ecological functioning and resilience.

The weighting scheme may be adjusted for specialized applications, provided scientific justification is documented.

3.8. Scoring Methodology

Each indicator is scored using a standardized five-point scale. Exceptional: 5; High: 4; Moderate:3; Low: 2; Very Low: 1. Scores are normalized according to indicator weights. The composite PRGI score is calculated as:

$$PRGI = \sum_{j=1}^4 \left(\frac{S_j}{5} \times 100 \right) W_j$$

S_j = average indicator score (1–5) for PRGI pillar j

W_j = fractional pillar weight (Hydrology 0.30, Biodiversity 0.30, Uniqueness 0.20, Ecosystem Services 0.20).

Or equivalently:

$$PRGI = 20 \sum_{j=1}^4 (S_j W_j)$$

The index is expressed on a 100-point scale. With indicator scores restricted to 1–5 and fractional pillar weights summing to 1, its attainable range is 20–100. Missing evidence is recorded as not assessed and is not assigned a zero score.

Every individual pillar may have its own sub indicators. For example, suppose Hydrology has four representative sub indicators, namely: **Natural Flow Regime, Perenniality, Recharge, and Floodplain Interaction**. The average indicator score may be calculated as shown in following Table.

Table 3.2: An Example of Calculation of Average Indicator Score of a Pillar

Hydrology Indicator	Score (1–5)
Natural flow regime	5
Perenniality	5
Recharge function	4
Floodplain interaction	4
Average indicator score	4.50

The normalized Hydrology pillar score is $4.50 \div 5 \times 100 = 90$
its weighted contribution is $90 \times 0.30 = 27.0$:

$$H = \frac{4.50}{5} \times 100 = 90$$

Thus, the original 1–5 indicator scores can be converted into a normalized pillar score of 90/100.

Its contribution to PRGI is then: $90 \times 0.30 = 27.0$

3.9. PRGI Grades

The composite score is interpreted using five ecological grades.

Table 3.3: Ecological Grades for Rivers and River Reaches

Grade	Score	Interpretation
Tier I	$85 \leq \text{PRGI} \leq 100$	Exceptional Ecological Significance
Tier II	$70 \leq \text{PRGI} < 85$	High Ecological Significance
Tier III	$55 \leq \text{PRGI} < 70$	Moderate Ecological Significance
Tier IV	$40 \leq \text{PRGI} < 55$	Limited Ecological Significance
Tier V	$20 \leq \text{PRGI} < 40$	Relatively Low Ecological Significance

These grades indicate ecological importance only. They should never be interpreted as measures of river degradation.

3.10. Interpretation of PRGI

The ecological grades guide conservation planning rather than restoration priorities.

Tier I: Represents rivers or rivers reaches of outstanding ecological significance possessing exceptional biodiversity, hydrological importance, uniqueness, and ecosystem services.

Management objective: Strict protection and long-term conservation.

Tier II: Represents rivers of high ecological significance supporting important ecological functions and biodiversity.

Management objective: Protection with carefully regulated sustainable use.

Tier III: Represents rivers of moderate ecological importance.

Management objective: Integrated conservation and sustainable management.

Tier IV: Represents River systems having relatively limited ecological significance but still contributing valuable local ecosystem functions.

Management objective: Maintain ecological functionality while supporting multiple uses.

Tier V: Represents rivers having comparatively lower intrinsic ecological significance.

Management objective: Appropriate management according to basin priorities while maintaining minimum ecological integrity.

Importantly, a Tier V River may still possess excellent functional health if natural processes remain intact.

3.11. Applications

PRGI provides valuable information for numerous river basin management activities. Major applications include:

- identification of ecologically important rivers
- conservation prioritization
- designation of protected river corridors
- biodiversity management
- environmental impact assessment
- ecological zoning
- river-sensitive infrastructure planning
- basin-scale conservation strategies
- ecological compensation planning
- national river inventories

PRGI also supports international commitments relating to biodiversity conservation and ecosystem restoration.

3.12. Relationship with Other Assessment Frameworks

Within the integrated framework developed in this guideline:

River Classification identifies **"What kind of river is this?"**, PRGI determines **"How important is this river under natural conditions?"**, River Health Assessment (RHA) evaluates **"How well is this river functioning at any instance?"**, Local Water Systems assessment is to determine **"How effectively do associated water systems contribute to basin functioning?"**

Each framework addresses a distinct question, ensuring that assessments remain complementary and free from conceptual overlap.

3.13. Limitations

PRGI should not be used as:

- a water quality index,
- a pollution assessment tool,
- a restoration priority index,
- an environmental compliance indicator, or
- a measure of present ecological health.

Instead, it quantifies the **intrinsic conservation value** of rivers under natural conditions.

3.14. Concluding Remarks

The Pristine River Grading Index provides a scientifically robust methodology for identifying and ranking the intrinsic ecological significance of rivers. By focusing on natural value rather

than present condition, PRGI establishes a strong foundation for conservation planning and complements the River Classification Framework described in the previous chapter.

Together, River Classification and PRGI answer the first two fundamental questions of integrated river assessment: **What type of river is it?** and **how important is it in its natural state?**

4. River Health Assessment (RHA)

4.1. Introduction

A river is far more than a flowing body of water. It is a dynamic, self-organizing geologically formed ecological system in which water, sediments, channel morphology, floodplains, riparian vegetation, aquatic organisms, groundwater, nutrients, and human interactions continuously influence one another. The long-term sustainability of a river depends not merely on the quality of its water but on the integrity of the physical, chemical, biological, and ecological processes that enable the river to function as a living system.

Traditional river assessments have frequently relied on water quality parameters as the primary indicator of river condition. While water quality is undoubtedly important, it represents only one manifestation of river functioning. A river may satisfy prescribed water quality standards while simultaneously suffering from disrupted sediment transport, fragmented habitats, altered flow regimes, disconnected floodplains, invasive species, or severely modified channel morphology. Conversely, a naturally sediment-rich Himalayan River may appear turbid but remain ecologically intact and highly functional.

Recognizing these limitations, the **River Health Assessment (RHA)** has been developed as an integrated, process-based framework for evaluating the present condition of rivers relative to their natural reference state.

Unlike conventional environmental indices, RHA evaluates how effectively the essential functions of a river continue to operate. It considers hydrological processes, channel and sediment dynamics, hydrochemistry, ecological integrity, habitat quality, longitudinal and lateral connectivity, and ecosystem services as interconnected dimensions of river health. By assessing these components together, RHA provides a holistic understanding of river condition and directly supports river restoration, environmental flow assessment, and integrated river basin management.

4.2. Purpose and Objectives

The River Health Assessment is designed to answer a single, clearly defined question: "**How well is the river functioning at any instance relative to its natural reference condition?**"

The principal objectives of RHA are:

- To evaluate the present functional condition of river reaches.
- To distinguish natural variability from anthropogenic degradation.
- To identify the principal drivers of ecological impairment.
- To support river restoration and rejuvenation planning.
- To guide environmental flow assessments.

- To prioritize management interventions.
- To monitor long-term changes in river condition.
- To provide a nationally consistent assessment framework applicable across diverse river systems in India.

Unlike PRGI, which evaluates ecological significance, RHA assesses current functionality. The two frameworks are therefore complementary rather than overlapping.

4.3. Conceptual Framework

The conceptual basis of RHA is founded on the understanding that **river health is the capacity of a river to sustain its characteristic physical, chemical, biological, and ecological functions over time while maintaining resilience to natural and anthropogenic disturbances.**

This definition emphasizes that rivers are not static entities, but dynamic systems maintained by interacting processes. Healthy rivers continuously transport water and sediment, exchange nutrients, sustain habitats, support biodiversity, interact with floodplains and groundwater, and deliver ecosystem services. Degradation occurs when one or more of these processes are impaired to the extent that the river can no longer perform its natural functions effectively.

Accordingly, RHA adopts a **functional systems approach** rather than a parameter-based approach. Instead of asking whether a river meets predefined standards for individual variables, RHA evaluates whether the river continues to function as an integrated ecological system.

4.4. Guiding Principles

The River Health Assessment is based on the following principles:

Reference Condition

Assessment shall always be undertaken relative to an appropriate reference condition representing the natural or least-disturbed state of the same river type. This ensures that naturally sediment-rich, mineralized, or seasonally variable rivers are evaluated according to their inherent characteristics rather than against universal standards.

Process-Based Assessment

River health depends on the integrity of interacting hydrological, geomorphological, hydrochemical, ecological, and biophysical processes. RHA therefore evaluates processes rather than isolated environmental parameters.

Reach-Based Assessment

River characteristics vary longitudinally. Consequently, assessments shall be conducted at the river reach level, with reach boundaries defined by major geomorphic, hydrological, or land-use transitions.

Integrated Assessment

Hydrology, sediment transport, habitat, ecology, connectivity, hydrochemistry, and ecosystem services are evaluated as mutually dependent components of a single river system.

Reference Condition

Data Availability and Transparency, allowing application using detailed field surveys, remote sensing, existing monitoring programmes, or expert assessments, provided that the methodology remains transparent and scientifically defensible.

4.5. Functional Dimensions of River Health

RHA is scored through six components: Hydrological Condition (H), Geomorphological Condition (G), Sediment Regime (S), Water and Environmental Quality (W), Habitat and Biodiversity (B), and Connectivity (C). The functional themes described in Section 4.5.1 – Section 4.5.8 explain the processes and outcomes informing these components; they are not eight separately weighted scores. Ecological integrity and habitat quality are assessed together under B. Ecosystem services are interpreted as cross-cutting outcomes and supporting evidence, without an additional composite weight.




Figure 4.1: Schematic Representation of Interrelated Functional Dimensions of River Health

4.5.1. Hydrological Function

Hydrology governs the quantity, timing, duration, frequency, and variability of river flows. It determines habitat availability, sediment transport, floodplain inundation, groundwater recharge, ecological connectivity, and ecosystem productivity. Key indicators include:

- Environmental flow regime
- Seasonal flow variability
- Flow permanence

- Flood pulse characteristics
- Groundwater interaction
- Degree of flow regulation
- Water abstraction pressure

4.5.2. Geomorphological Function

Geomorphology reflects the capacity of the river to maintain its characteristic channel form and floodplain processes. Assessment considers:

- Channel stability
- Channel pattern
- Bank condition
- Floodplain integrity
- Channel migration
- Natural geomorphic processes

4.5.3. Sediment Continuity

Sediment is fundamental to river functioning. It shapes channels, forms habitats, sustains deltas, replenishes floodplains, and supports ecological processes. Indicators include:

- Sediment supply
- Sediment transport continuity
- Sediment balance
- Bed material condition
- Reservoir trapping effects
- Channel incision or aggradation

4.5.4. Hydrochemical Function

Hydrochemistry influences nutrient cycling, aquatic productivity, ecological resilience, and overall river metabolism. Assessment considers:

- Dissolved oxygen
- Nutrient balance
- Organic pollution
- Salinity
- pH
- Electrical conductivity
- Toxic contaminants

Importantly, naturally occurring hydrochemical characteristics should be distinguished from pollution-induced changes.

4.5.5. Ecological Integrity

Ecological integrity represents the capacity of the river to sustain native biological communities and ecological interactions. Representative indicators include:

- Native fish assemblages

- Macroinvertebrate diversity
- Riparian vegetation
- Aquatic biodiversity
- Invasive species
- Trophic structure
- Ecological resilience

4.5.6. Habitat Quality

Habitat quality reflects the availability and diversity of physical environments supporting aquatic and riparian organisms. Assessment includes:

- Habitat diversity
- Instream woody material
- Pools and riffles
- Riparian condition
- Substrate diversity
- Refuge habitats
- Wetland connectivity

4.5.7. Connectivity

Connectivity enables the movement of water, sediments, nutrients, organisms, and energy throughout river systems. RHA evaluates:

- Longitudinal connectivity
- Lateral connectivity with floodplains
- Vertical connectivity with groundwater
- Tributary connectivity
- Fish passage
- Wetland connectivity

4.5.8. Ecosystem Services

Healthy rivers provide numerous services supporting both ecosystems and society. Assessment considers:

- Groundwater recharge
- Flood attenuation
- Natural purification
- Fisheries
- Recreation
- Cultural values
- Climate regulation
- Carbon sequestration
- Livelihood support

4.6. Integrated Functional Assessment Framework

The six RHA components collectively define present river-health condition. The functional themes in Section 4.5 provide process-based evidence for these components. Disruption of sediment continuity may alter channel morphology, reduce habitat diversity, disconnect floodplains, impair biodiversity and diminish ecosystem services. Assess these linked effects through H, G, S, W, B and C, assign each indicator to a declared component, and prevent duplicate weighting of the same evidence.

4.7. Assessment Indicators

Table 4.1 presents the matrix of River Health Assessment (RHA) outlining Functional Dimensions, Objectives, Indicators, Assessment Considerations along with Typical Stressors. This table will form the operational basis for RHA implementation and can be adapted to basin-specific conditions while maintaining national consistency.

4.8. RHA Assessment Indicators and Weighting Scheme

RHA integrates six component scores using Table 4.2: H 20%, G 15%, S 15%, W 20%, B 20% and C 10%. The functional themes described in Section 4.5 supply evidence for these six components and do not receive additional independent weights.

The recommended weighting emphasizes the primary drivers of river functionality—hydrology, geomorphology, sediment continuity, and ecology—while ensuring that hydrochemistry, habitat quality, connectivity, and ecosystem services are adequately represented.

Table 4.1: River Health Assessment (RHA): Functional Dimensions, Objectives, Indicators and Assessment Considerations

Functional Theme	Assessment Objective	Representative Indicators	Typical Data Sources / Assessment Methods	Typical Stressors
Hydrological Function	Evaluate the extent to which the river maintains its natural flow regime and hydrological processes.	<ul style="list-style-type: none"> ➤ Environmental flow compliance ➤ Seasonal flow variability ➤ Baseflow contribution ➤ Flood pulse characteristics ➤ Flow permanence ➤ Water abstraction intensity ➤ Groundwater–surface water interaction 	Hydrological records, flow gauging stations, environmental flow studies, groundwater monitoring, remote sensing, hydrological models	Dams, barrages, diversions, groundwater abstraction
Geomorphological Function	Assess the ability of the river to maintain natural channel form, floodplain processes and geomorphic adjustments.	<ul style="list-style-type: none"> ➤ Channel stability ➤ Bank erosion/accretion ➤ Channel pattern ➤ Floodplain connectivity ➤ Channel migration ➤ Bedform diversity ➤ River Styles condition 	Topographic surveys, satellite imagery, River Styles assessment, geomorphic mapping, field surveys	Channelization, embankments, sand mining
Sediment Continuity	Determine whether sediment production, transport and deposition processes remain functionally connected.	<ul style="list-style-type: none"> ➤ Sediment load ➤ Sediment continuity ➤ Bed material composition ➤ Reservoir trapping effects ➤ Channel incision/aggradation ➤ Sediment balance 	Sediment monitoring, bathymetric surveys, reservoir sedimentation studies, geomorphic investigations	Dams, excessive mining, bank protection

Table 4.1 Continued on next page

Functional Dimension	Assessment Objective	Representative Indicators	Typical Data Sources / Assessment Methods	Typical Stressors
Hydrochemical Function	Assess whether natural water chemistry supports ecological functioning.	<ul style="list-style-type: none"> ➤ Dissolved oxygen ➤ Biochemical Oxygen Demand (BOD) ➤ Nutrients (N & P) ➤ pH ➤ Electrical conductivity ➤ Salinity ➤ Toxic contaminants ➤ Water temperature 	Water quality monitoring programmes, laboratory analysis, pollution inventories	Sewage, industrial discharge, agricultural runoff
Ecological Integrity	Evaluate the ability of the river to sustain native biological communities and ecological processes.	<ul style="list-style-type: none"> ➤ Native fish diversity ➤ Macroinvertebrate communities ➤ Riparian vegetation ➤ Aquatic flora ➤ Invasive species ➤ Species richness ➤ Ecological resilience 	Biodiversity surveys, ecological monitoring, fisheries assessments, vegetation mapping	Habitat loss, invasive species, overfishing
Habitat Quality	Assess the diversity, quality and availability of aquatic and riparian habitats.	<ul style="list-style-type: none"> ➤ Habitat heterogeneity ➤ Pools, riffles and runs ➤ Substrate diversity ➤ Large woody debris ➤ Riparian buffer condition ➤ Wetland association ➤ Refuge habitats 	Habitat surveys, River Habitat Survey (RHS), drone imagery, field observations	Encroachment, riparian clearing, channel modification

Table 4.1 Continued on next page

Functional Dimension	Assessment Objective	Representative Indicators	Typical Data Sources / Assessment Methods	Typical Stressors
Connectivity	Evaluate the continuity of ecological and hydrological linkages within the river corridor.	<ul style="list-style-type: none"> ➤ Longitudinal connectivity ➤ Lateral floodplain connectivity ➤ Vertical groundwater connectivity ➤ Tributary connectivity ➤ Fish passage ➤ Wetland connectivity 	GIS analysis, dam inventories, floodplain mapping, field assessment, ecological connectivity studies	Weirs, barrages, culverts, levees
Ecosystem Services	Assess the river's capacity to provide ecological and socio-economic benefits.	<ul style="list-style-type: none"> ➤ Groundwater recharge ➤ Flood attenuation ➤ Nutrient cycling ➤ Carbon sequestration ➤ Fisheries ➤ Recreation ➤ Cultural and spiritual values ➤ Livelihood support 	Ecosystem service valuation, socio-economic surveys, land-use analysis, stakeholder consultation	Urbanization, land-use change, over-extraction

Assessment Notes

1. Each functional dimension should be evaluated relative to an appropriate **Reference Condition** established for that river type.
2. Indicators may be refined according to basin characteristics while retaining the six scored RHA components H, G, S, W, B and C.
3. Quantitative data should be used wherever available. Where data are limited, structured expert judgement supported by field observations may be adopted.
4. Map the functional themes to scored components as follows: **Hydrological Function → H; Geomorphological Function → G; Sediment Continuity → S; Hydro-chemical Function → W; Ecological Integrity and Habitat Quality → B; Connectivity → C**. Interpret Ecosystem Services as cross-cutting outcomes and supporting evidence, without a separate weight. Aggregate declared indicators into their assigned component scores and apply the six component weights in Table 4.2.
5. The framework is designed for application at the **river reach** scale, recognizing that functional characteristics vary longitudinally and laterally.

Table 4.2: RHA Components and Weightages

Functional Dimension	Weight (%)	Rationale
Hydrological Condition (H)	20	Primary driver of river processes
Geomorphological Condition (G)	15	Controls channel stability and evolution
Sediment Regime (S)	15	Maintains channel form and habitats
Water & Environmental Quality (W)	20	Reflects biological functioning and resilience
Habitat & Biodiversity (B)	20	Sustains aquatic and riparian biodiversity
Connectivity (C)	10	Retain the weights H 20%, G 15%, S 15%, W 20%, B 20% and C 10%

The weighting scheme may be modified for specific river basins where scientific evidence demonstrates the need for regional adjustments. However, any modification should preserve the integrated nature of the framework.

4.9. Assessment Indicators

Each of the six RHA components is evaluated using declared measurable indicators selected from the relevant functional themes. Selection of indicators should consider scientific validity, data availability, reproducibility, and applicability across diverse river systems.

Table 4.3: Functional Themes and Representative Indicators

Functional Theme	Representative Indicators
Hydrology	Environmental flows, flow variability, abstraction, flood pulses, groundwater interaction
Geomorphology	Channel stability, bank erosion, floodplain integrity, channel pattern, geomorphic diversity
Sediment Continuity	Sediment balance, sediment transport continuity, bed material, reservoir trapping
Hydrochemistry	Dissolved oxygen, nutrients, organic pollution, conductivity, pH, toxic substances
Ecological Integrity	Fish diversity, macroinvertebrates, riparian vegetation, invasive species, aquatic biodiversity
Habitat Quality	Habitat diversity, substrate composition, pools, riffles, woody debris, refuge habitats
Connectivity	Longitudinal, lateral, vertical and tributary connectivity; fish passage
Ecosystem Services	Flood attenuation, groundwater recharge, fisheries, recreation, cultural value

These indicators support the six-component RHA framework. Use the theme-to-component mapping in the notes to Table 4.1, combine Ecological Integrity and Habitat Quality within B, and record Ecosystem Services as outcomes or supporting evidence without a separate composite weight. Additional basin-specific indicators require a documented rationale.

4.10. Scoring Methodology

Each indicator is evaluated using a five-point scoring scale based on comparison with the established reference condition.

Table 4.4: Indicator Scoring Scale

Score	Functional Condition
5	Pristine/Near-reference
4	Slightly Modified
3	Moderately Altered
2	Heavily Altered
1	Critically Degraded

Scores should be assigned using quantitative data wherever available. Where quantitative information is lacking, structured expert judgement supported by field observations may be used.

For each functional dimension,

$$\text{Dimension Score} = \frac{\sum(\text{IndicatorScore} \times \text{IndicatorWeight})}{\sum \text{IndicatorWeight}}$$

The overall RHA Score is computed as:

$$RHA = 0.20H + 0.15G + 0.15S + 0.20W + 0.20B + 0.10C$$

H, G, S, W, B and C are component scores on 1-5 scale, each calculated from its declared indicator scores and normalized indicator weights. The RHA Score ranges from 1.0 to 5.0. Assign the RHA Class from the unrounded score and round only the displayed result.

4.11. River Health Classes

The River Health Score is interpreted using five Health Classes.

Table 4.5: River Health Classes

Class	RHS Score	Functional Status
Class A	$4.50 \leq RHA \leq 5.00$	Pristine/Near-reference
Class B	$3.50 \leq RHA < 4.50$	Slightly Modified
Class C	$2.50 \leq RHA < 3.50$	Moderately Altered
Class D	$1.50 \leq RHA < 2.50$	Heavily Altered
Class E	$1.00 \leq RHA < 1.50$	Critically Degraded

These classes provide a nationally consistent framework while allowing river-type-specific interpretation. The RHA assessment classes are schematically presented in Figure 4.2.




Figure 4.2: River Health Assessment Classes

4.12. Interpretation of Health Classes

Class A – Pristine/Near-reference: The river retains almost all natural hydrological, geomorphological, ecological, and biogeochemical functions. Habitat diversity is high, ecological connectivity remains intact, and ecosystem services are fully maintained.

Management Objective: *Protect existing conditions through conservation and preventive management.*

Class B – Slightly Modified: Minor alterations have occurred, but the river continues to function effectively with minimal impairment of ecological processes. Examples include limited flow regulation, localized bank stabilization, or moderate land-use change.

Management Objective: *Maintain existing condition while preventing further degradation.*

Class C – Moderately Altered: Noticeable degradation has occurred in several functional dimensions. Examples include:

- altered flow regime
- fragmented habitats
- reduced biodiversity
- moderate pollution
- interrupted sediment transport

However, ecological recovery remains feasible.

Management Objective: *Active restoration and adaptive management.*

Class D – Heavily Altered: Multiple River processes have been substantially impaired. Typical characteristics include:

- major hydrological alteration
- extensive channel modification
- severe pollution
- disrupted connectivity
- degraded habitats
- declining biodiversity

Management Objective: *Comprehensive rehabilitation supported by basin-wide interventions.*

Class E – Critically Degraded: River functioning has largely collapsed. Hydrological, geomorphological, ecological and biogeochemical processes are severely impaired. Examples include:

- heavily urbanized channels
- concrete-lined rivers
- severely polluted reaches
- disconnected floodplains
- ecological collapse

Management Objective: *Long-term ecological recovery through integrated river restoration programmes.*

4.13. Management Decision Framework

RHA is intended not merely as an assessment tool but as a decision-support framework.

Table 4.6: Functional Health Classes and Recommended Management Response

Functional Health Class	Recommended Management Strategy
Class A	Long-term protection and conservation
Class B	Preventive management and monitoring
Class C	Targeted ecological restoration
Class D	Comprehensive river rehabilitation
Class E	Intensive basin-scale restoration and recovery

This linkage between assessment and management enables direct application in river basin planning.

4.14. Integration with River Classification and PRGI

The three principal assessment frameworks developed in this guideline are complementary and sequential.

River Classification answers **What type of river is this?** It establishes the natural physical identity of the river.

Pristine River Grading Index (PRGI) answers **How ecologically important is this river under natural conditions?** It identifies intrinsic conservation value.

River Health Assessment (RHA) answers **How well is this river functioning presently?** It evaluates present ecological condition relative to the river's natural reference state.

RCF, PRGI and RHA constitute the three main-river assessment layers. FACLWS provides the complementary functional assessment and diagnosis of associated Local Water Systems.

4.15. Applications

RHA has broad applicability across river basin planning and management. Major applications include:

- River rejuvenation programmes
- Environmental flow assessment
- Basin restoration planning
- Conservation prioritization
- Climate resilience assessment
- Wetland restoration
- Dam operation and sediment management
- Urban river restoration
- Monitoring of river health trends
- Adaptive basin management
- Evaluation of restoration outcomes
- Reporting under national river conservation programmes

The framework is equally applicable to Himalayan, alluvial, peninsular, coastal, spring-fed, and ephemeral river systems, provided assessments are undertaken relative to appropriate reference conditions.

4.16. Strengths and Limitations

Strengths

The RHA framework:

- adopts a process-based rather than parameter-based approach;
- integrates physical, chemical, biological, and ecological dimensions into a single assessment;
- distinguishes natural variability from anthropogenic degradation;
- is compatible with River Styles, environmental flow assessment, and integrated river basin management;
- provides direct guidance for restoration and conservation planning; and
- is scalable from individual reaches to entire river basins.

Limitations

Like any assessment framework, RHA depends on the availability and quality of data. In data-scarce regions, expert judgement may be required, and uncertainties should be documented transparently. The framework is therefore intended to support adaptive management, with periodic reassessment as improved information becomes available.

4.17. Concluding Remarks

The River Health Assessment constitutes the central operational component of the integrated river assessment framework presented in this guideline. RHA provides a more coherent, scientifically robust, and management-oriented assessment of river condition.

When used together with the **River Classification Framework** (Chapter 2) and the **Pristine River Grading Index** (Chapter 3), RHA completes the three-tier assessment hierarchy:

- **River Classification** defines the river's intrinsic identity.
- **PRGI** determines its intrinsic ecological significance.
- **RHA** evaluates its present condition relative to its natural reference state.

Together, River Classification, PRGI and RHA provide the three complementary main-river assessment layers: River Classification establishes identity and reference context; PRGI establishes intrinsic ecological significance; and RHA establishes present river-health condition. FACLWS extends the diagnosis to associated Local Water Systems. This integrated hierarchy enables managers to distinguish whether a river should be **protected, maintained, restored, or rehabilitated**, thereby providing a sound scientific basis for river basin planning under the **Ganga River Basin Management Plan (GRBMP 2.0)** and for application across diverse river systems in India.

5. Functional Assessment and Classification of Local Water Systems (FACLWS)

5.1. Introduction

Healthy river basins comprise not only the main river channel but also an extensive network of interconnected local water systems, including first- and second-order streams, natural drains, floodplain channels, wetlands, ponds, lakes, springs, reservoirs, and urban drainage systems. These systems collectively regulate hydrological connectivity, groundwater recharge, sediment transport, nutrient cycling, habitat diversity, flood attenuation, and ecological resilience. Their cumulative influence often determines the functional condition of the river itself.

FACLWS is a diagnostic framework. It identifies the Local Water System, establishes its natural/intended functions, assesses its present functional condition, identifies functional deficits and determines its influence on the receiving river. Planning, intervention and long-term management are undertaken subsequently through PMLWS.

Conventional river management has generally focused on the main river channel, while associated local water systems have been managed independently under different institutional and administrative frameworks. Urban drains are treated primarily as wastewater conveyance systems, ponds as isolated water bodies, wetlands as land-use categories, and natural streams as local drainage features. Such fragmented approaches fail to recognize that these systems collectively constitute the functional infrastructure of a river basin.

Under the **Ganga River Basin Management Plan (GRBMP 2.0)**, basin management extends beyond the river channel to encompass the entire aquatic network. Accordingly, FACLWS constitutes the fourth assessment and diagnostic component of IRAF. Its assessment units are individual Local Water Systems and their connected networks; its findings inform subsequent PMLWS planning and management. Unlike the preceding chapters, which focus on the intrinsic characteristics and functional health of rivers, this chapter evaluates the **functional performance of local water systems** in sustaining basin-scale ecological and hydrological processes.

The emphasis is therefore not on restoring every local water body to pristine conditions but on maximizing its contribution to river basin functioning through appropriate conservation, restoration, rehabilitation, or adaptive management. The schematic representation of various treatment systems for improving quality of used- water/ sewage/wastewater/ such as electro-mechanical (STPs/ETPs) systems, nature-based systems and natural systems for balancing

utilization of land, energy and human resources and developing local ecological infrastructure is schematically presented in Figure 5.1.




Figure 5.1: Schematic Representation of Various Treatment Systems for Used Water: Electro-Mechanical Systems, Nature Based Systems and Natural Systems for Balancing Utilization of Land, Energy and Human Resources and Developing Local Ecological Infrastructure

5.2. Need for Utilizing Local Water System to Develop Ecological Infrastructure

Local water systems vary enormously in their origin, morphology, hydrology, ecological characteristics, and present condition. Some are entirely natural, while others have evolved into highly modified but functionally important systems. For example:

- A natural spring-fed stream may possess exceptional ecological value.
- An urban drain may no longer retain natural characteristics but may significantly improve river health if transformed into an ecologically functioning treatment corridor.
- A restored wetland may provide flood attenuation, nutrient removal, groundwater recharge, and habitat creation despite substantial historical modification.
- A village pond may contribute to local groundwater recharge and biodiversity even though it has limited connection to the main river.

Assessment based solely on naturalness is therefore inadequate for these systems. Instead, their contribution to basin functioning should be evaluated according to the ecological, hydrological, treatment, habitat, and socio-cultural functions they presently perform or can realistically perform following restoration.

5.3. Objectives

The assessment of Local Water Systems seeks to:

- Evaluate the contribution of local water systems to river basin functioning.
- Recognize both natural and restored systems according to their functional performance.
- Support decentralized water management and nature-based solutions.
- Guide restoration prioritization.
- Strengthen hydrological and ecological connectivity within river basins.
- Promote sustainable urban and rural water management.
- Complement the River Health Improvement efforts.

5.4. Guiding Principles

The framework is based on six principles.

Functional Performance: Assessment emphasizes how effectively a system performs ecological and hydrological functions rather than how closely it resembles its original natural condition.

Basin Connectivity: Every local water system should be evaluated in relation to its contribution to the broader river basin.

Multiple Functions: Most systems simultaneously perform hydrological, ecological, water quality, habitat, and socio-economic functions.

Context Sensitivity: Functional expectations should reflect climatic, geomorphological, and land-use contexts.

Restoration Potential: Modified systems should be assessed according to achievable ecological performance rather than unrealistic restoration targets.

Adaptive Management: Assessment should support continuous improvement through monitoring and periodic reassessment.

5.5. Local Water Systems

Local Water Systems are assessed through five functional dimensions: Hydrological; Water-quality; Ecological; Geomorphic/sediment; and Human-support functions, as shown in Figure 5.2.




Figure 5.2: Functions of Local Water Systems

The framework recognizes six principal categories of local water systems.

5.5.1. Natural Streams

Small perennial or seasonal streams forming the upper drainage network. Primary functions include:

- flow generation
- sediment transport
- habitat provision
- groundwater interaction
- ecological connectivity

5.5.2. Drains

Artificial or modified drainage channels conveying stormwater, wastewater, or mixed flows. Although often degraded, properly managed drains can perform valuable functions including:

- pollutant attenuation
- decentralized wastewater treatment
- flood conveyance
- ecological corridors
- groundwater recharge (where appropriate)

5.5.3. Wetlands

Natural or constructed wetlands perform critical ecological functions including:

- nutrient removal
- flood storage
- habitat support

- carbon sequestration
- groundwater recharge
- biodiversity conservation

Wetlands should therefore be regarded as integral components of river systems rather than isolated water bodies.

5.5.4. Ponds and Lakes

Village ponds, urban ponds, lakes, oxbow lakes, and reservoirs contribute to:

- local storage
- groundwater recharge
- biodiversity
- fisheries
- recreation
- landscape amenity

Their functional value depends more upon ecological performance than size.

5.5.5. Floodplain Water Bodies

Floodplain lakes, side channels, abandoned meanders, and seasonal depressions provide:

- flood attenuation
- sediment storage
- fish breeding
- nutrient cycling
- biodiversity refugia

Maintaining connectivity with the river is essential for sustaining these functions.

5.5.6. Nature-Based Treatment Systems

Constructed wetlands, polishing ponds, decentralized treatment systems, ecological treatment corridors, and vegetated drains increasingly form part of integrated urban water management. Although engineered, these systems contribute substantially to river basin health through:

- wastewater polishing
- nutrient removal
- habitat creation
- landscape enhancement
- groundwater recharge
- climate resilience

5.6. Functional Dimensions

Evaluation of each local water system may be done using five functional dimensions and the sequential diagnostic pathway (Figure 5.3).

Hydrological Functions: Flow regulation, storage, recharge/discharge, flood attenuation and hydrological connectivity.

Water-Quality Functions: Pollutant attenuation, nutrient transformation, organic-matter processing and compatible natural or engineered treatment.

Ecological Functions: Native biodiversity, aquatic and riparian habitat, ecological processes, refugia and ecological connectivity.

Geomorphic/sediment Functions: Sediment storage and transfer, side-channel and floodplain morphology, energy dissipation and erosion–deposition processes.

Human-Support Functions: compatible water use, livelihoods, flood-risk reduction, recreation, education, landscape amenity and cultural values. Assess each dimension against the system’s natural or intended role. Functions that are not applicable are identified explicitly; every system is not expected to perform all functions equally.




Figure 5.3: Schematic Representation of the Five Functional Dimensions of the Local Water System (top figure) and Sequential Diagnostic Pathway for Functional Assessment and Classification of Local Water Systems (FACLWS) (bottom figure)

5.7. Assessment Indicators

Table 5.1: Functional Assessment Framework

Functional Dimension	Representative Indicators
Hydrological Functions	Storage, recharge, flood attenuation, flow regulation
Ecological Functions	Biodiversity, vegetation, ecological integrity
Water-Quality Functions	Pollutant removal, nutrient attenuation, self-purification
Geomorphic Functions	Habitat diversity, connectivity, refuge value
Human-Support Functions	Recreation, aesthetics, education, cultural importance

5.8. Functional Performance Categories

Local water systems are classified according to their functional performance.

Table 5.2: Functional Categories

Category	Functional Status	Interpretation
A	Highly Functional	Performs most ecological and hydrological functions effectively
B	Functionally Adequate	Performs major functions with moderate limitations
C	Functionally Impaired	Significant reduction in functional performance
D	Non-functional	Performs few ecological or hydrological functions

FACLWS Categories A–D describe the functional performance of associated Local Water Systems. They are distinct from PRGI Tiers and RHA Classes and are assigned from the documented functional evidence; this table does not prescribe numerical cut-offs.

5.9. Diagnostic Requirements for PMLWS

Category A: Category A identifies functions requiring continued protection and monitoring emphasis:

- conservation
- preventive maintenance
- ecological monitoring

Category B: Category B identifies specific functional limitations and their causes:

- habitat enhancement
- connectivity improvement
- pollution reduction

Category C: Category C identifies substantial functional deficits, pressures, connectivity constraints and recovery requirements:

- ecological restoration
- flow improvement
- sediment management
- decentralized treatment

Category D: Category D identifies functions that are absent or effectively lost, together with constraints, risks and the evidence required to determine a feasible future role. FACLWS records these management requirements. Subsequent PMLWS planning selects and prioritises protection, sustenance, restoration, rehabilitation, reconnection, optimisation/repurposing or justified augmentation measures and prepares the management plan:

- nature-based restoration
- ecological engineering
- integrated urban water management
- basin-scale interventions

5.10. Functional Assessment Matrix

Table 5.3: FACLWS Functional Diagnosis Record

Functional Dimension	Expected Function and Reference	Observed Performance and Evidence	Functional Deficit and River Influence	Confidence
Hydrological functions				
Water-quality functions				
Ecological functions				
Geomorphic/sediment functions				
Human-support functions				

Report the five-dimension functional profile and the evidence supporting the Category A–D interpretation in Table 5.2. Report a composite score only when a system-specific schedule declares the applicable indicators, reference expectations, common scoring scale, normalized weights, treatment of non-applicable indicators and numerical category thresholds before scoring. Calculate the composite as the sum of each dimension score multiplied by its declared fractional weight; the applicable weights sum to 1. Report the dimension profile, weights and thresholds with the result. Until this schedule is complete, enter ‘Not calculated’ for the composite score.

5.11. Integration with RHA

The Functional Classification of Local Water Systems complements the River Health Assessment by evaluating the supporting aquatic infrastructure of the river basin. The relationship between the two frameworks may be summarized as follows:

Table 5.4: Relationship between RHA and FACLWS

River Health Assessment (RHA)	Functional Assessment and Classification of Local Water Systems (FACLWS)
Main river channel	Streams, wetlands, ponds, drains and associated water bodies
Evaluates river functioning	Evaluates supporting system performance
River-scale assessment	Local system assessment
Diagnosis of present river-health condition and reach-management requirements	Diagnosis of local-system functions, deficits and management requirements for subsequent PMLWS planning
Basin health outcome	Basin health contributors

Together, they provide a comprehensive understanding of both the **condition of the river** and the **performance of the supporting aquatic network**.

5.12. Applications

The Functional Classification of Local Water Systems supports:

- Urban river rejuvenation
- Decentralized wastewater management
- Wetland restoration
- Floodplain management
- Blue-green infrastructure planning
- Climate adaptation
- Stormwater management
- Groundwater recharge programmes
- Smart Cities and AMRUT initiatives
- Basin-scale ecological restoration

It is particularly valuable in rapidly urbanizing catchments, where restoring the functional contribution of local water systems can significantly improve downstream river health.

5.13. Relationship with the Integrated Assessment Framework

The four assessment components introduced in this guideline now form a complete hierarchy:

Table 5.5: Hierarchy for Four Suggested Assessment Components in Integrated Assessment

Framework	Principal Question	Assessment Focus
River Classification	What type of river is it?	Intrinsic physical identity
Pristine River Grading Index (PRGI)	How ecologically important is it under natural conditions?	Intrinsic ecological significance
River Health Assessment (RHA)	How well is it functioning presently?	Present river condition

Framework	Principal Question	Assessment Focus
Functional Assessment and Classification of Local Water Systems (FACLWS) as Supporting Ecological Infrastructure in a River Basin and/or an Administrative Unit (District)	How effectively do associated water systems support basin functioning?	Functional performance of supporting aquatic systems

This hierarchy ensures that each framework addresses a distinct management question, avoiding conceptual overlap while supporting integrated river basin planning.

5.14. Concluding Remarks

Local water systems are indispensable components of healthy river basins. Their contribution extends beyond water storage and drainage to include groundwater recharge, flood moderation, nutrient cycling, biodiversity conservation, pollutant attenuation, habitat creation, and socio-cultural benefits. Assessing these systems solely based on naturalness overlooks their current and potential functional roles within the basin.

The **Functional Assessment and Classification of Local Water Systems (FACLWS)** provides a practical framework for evaluating these contributions and complements the River Classification, PRGI, and RHA by extending assessment beyond the main river channel to the broader aquatic network. Together, these four frameworks establish an integrated and scalable foundation for river basin planning, restoration, and sustainable water resources management under **GRBMP 2.0**.

6. Integration into River Basin Management

6.1. Introduction

The preceding chapters have presented four complementary assessment frameworks that characterize rivers from different but interconnected perspectives. The **River Classification Framework** establishes the intrinsic identity of river systems; the **Pristine River Grading Index (PRGI)** evaluates their intrinsic ecological significance; the **River Health Assessment (RHA)** assesses their present condition; and the **Functional Assessment and Classification of Local Water Systems (FACLWS)** evaluates the contribution of associated aquatic systems to basin-scale functioning.

Individually, each framework provides valuable scientific information. However, their greatest strength lies in their **integrated application** within the context of river basin management. River restoration, conservation, environmental flow assessment, urban river rejuvenation, and sustainable water resources management require decisions that simultaneously consider river identity, ecological significance, present condition, and the functioning of supporting aquatic systems.

Accordingly, this chapter presents an integrated decision-support framework that translates scientific assessment into management action. Rather than generating isolated indices, the four frameworks collectively guide **what should be protected, what should be restored, what should be rehabilitated, and what should be sustainably managed**.

6.2. From Assessment to Management

The integrated framework follows a logical sequence of decision-making.

Step 1 – Identify the River

Determine the intrinsic characteristics of the river using the River Classification Framework.

Step 2 – Determine Ecological Significance

Assess the intrinsic conservation value through PRGI.

Step 3 – Assess River Health

Evaluate the present functional condition using RHA.

Step 4 – Diagnose Associated Local Water Systems

Assess expected functions, present functional condition and deficits through FACLWS.

Step 5 – Integrated Diagnosis

Synthesise RCF, PRGI, RHA and FACLWS outputs to identify principal pressures, causes of impairment, functional deficits and management requirements.

Step 6 – Coordinated Planning and Management

Prepare distinct, coordinated Reach-Specific River Management and PMLWS plans.

Step 7 – Implementation and Monitoring

Implement the agreed measures and monitor their effects on river condition and local-system functions.

Step 8 – Reassessment and Adaptive Management

Use monitoring results to reassess diagnosis and revise both management plans.

This sequence ensures that management decisions are scientifically defensible and context specific.

6.3. Integrated River Assessment Hierarchy

RCF establishes identity and reference context; PRGI, RHA and FACLWS provide complementary assessments at their appropriate operational scales. Integrated Diagnosis combines these findings to inform two coordinated management pathways. Integrated Diagnosis combines these findings to inform two coordinated management pathways.




Figure 6.1: Integrated River Assessment Hierarchy

6.4. Decision Matrix for River Basin Management

The combined use of PRGI and RHA provides the basis for selecting management priorities.

Table 6.1: Integrated Decision Matrix for River Basin Management

PRGI	RHA	Management Priority
High	High	Strict conservation and protection
High	Moderate	Prevent further degradation and restore impaired functions
High	Low	Highest priority for ecological restoration
Moderate	High	Sustainable management and monitoring

PRGI	RHA	Management Priority
Moderate	Moderate	Targeted restoration and adaptive management
Moderate	Low	Basin rehabilitation with phased interventions
Low	High	Maintain ecosystem functions while supporting multiple uses
Low	Moderate	Improve selected ecological functions
Low	Low	Rehabilitation according to basin priorities

This matrix recognizes that ecological importance and river health are independent characteristics.

6.5. River Basin Zonation

The integrated framework supports basin-wide zoning based on ecological significance and functional condition.

Conservation Zones: River reaches exhibiting both high PRGI and high RHA should be designated as conservation zones where ecological integrity is maintained through preventive management. Examples include:

- pristine headwaters
- protected forest reaches
- ecologically sensitive tributaries

Restoration Zones: River reaches with high ecological significance, but poor river health requires active restoration. Typical interventions include:

- environmental flow restoration
- pollution control
- habitat enhancement
- sediment management
- floodplain reconnection

Sustainable Use Zones: River reaches with moderate ecological significance, and good river health may accommodate carefully managed water resource development while maintaining ecological functions.

Rehabilitation Zones: Highly degraded river reaches require integrated rehabilitation involving hydrological, geomorphological, ecological, and socio-economic interventions. Urban rivers frequently fall within this category.

6.6. Environmental Flow Management

- Environmental flows represent one of the most important applications of the integrated framework.
- River Classification identifies the natural hydrological regime.
- PRGI establishes ecological flow requirements for maintaining biodiversity and ecosystem services.

- RHA evaluates present hydrological functioning.
- FACLWS identifies opportunities for improving local hydrological processes through decentralized interventions and by utilizing Local Water Systems as Ecological Infrastructure for river rejuvenation and conservation.

Together, these frameworks support scientifically robust environmental flow assessments that extend beyond minimum flow prescriptions to include seasonal variability, flood pulses, sediment transport, ecological connectivity, and groundwater interactions.

6.7. River Restoration Planning

Effective river restoration should address the underlying processes responsible for degradation rather than merely improving water quality. **The integrated framework enables restoration planning through the following sequence:**

- Establish the natural river type.
- Identify the reference condition.
- Evaluate functional impairment.
- Diagnose dominant stressors.
- Prioritize restoration measures.
- Monitor functional recovery.

Typical restoration measures include:

- environmental flow restoration
- floodplain reconnection
- sediment continuity restoration
- riparian vegetation recovery
- wetland restoration
- habitat enhancement
- pollution reduction
- decentralized wastewater treatment
- barrier removal
- ecological engineering

6.8. Nature-Based Solutions

Nature-Based Solutions (NbS) form an essential component of basin-scale river management. Examples include:

- constructed wetlands
- ecological treatment corridors
- floodplain restoration
- riparian buffer establishment
- green stormwater infrastructure
- riverbank bioengineering

- recharge wetlands
- ecological ponds

These interventions improve both RHA and FACLWS scores while enhancing climate resilience and reducing infrastructure costs.

6.9. Urban River Rejuvenation

Urban rivers present unique management challenges due to altered hydrology, wastewater discharges, channel modification, encroachment, and loss of floodplain connectivity. **The integrated framework supports urban river rejuvenation through:**

- restoration of environmental flows
- decentralized wastewater management
- rehabilitation of urban drains as ecological corridors
- reconnecting floodplains where feasible
- restoration of ponds and wetlands
- enhancement of riparian green infrastructure
- community stewardship

Rather than attempting to recreate pristine conditions, urban restoration should focus on maximizing functional performance and ecosystem services.

6.10. Climate Change Adaptation

Climate change is expected to alter river hydrology through changing precipitation patterns, glacier retreat, increased evapotranspiration, extreme floods, prolonged droughts, and sea-level rise. **The integrated framework supports climate adaptation by:**

- protecting ecologically resilient river reaches;
- restoring natural hydrological processes;
- improving groundwater recharge;
- maintaining sediment continuity;
- enhancing floodplain storage;
- strengthening ecological connectivity; and
- increasing the adaptive capacity of local water systems.

Functional river systems are inherently more resilient to climate variability than highly modified systems.

6.11. Basin Governance

Successful implementation requires governance structures that recognize rivers as integrated socio-ecological systems rather than sectoral infrastructure. **Key governance principles include:**

- basin-scale planning;
- inter-agency coordination;
- adaptive management;
- stakeholder participation;
- science-based decision-making;
- periodic reassessment; and
- transparent monitoring.

The integrated assessment framework provides a common scientific language for engineers, ecologists, planners, policymakers, and local communities.

6.12. Monitoring and Adaptive Management

River systems continuously evolve in response to natural variability and human interventions. Consequently, management should follow an adaptive cycle (Figure 6.2):




Figure 6.2: Cyclic Process of Human-River Interaction and Adaptive Management of River System

Repeated RHA and FACLWS assessments, together with PMLWS performance monitoring, provide quantitative evidence of functional recovery, river-health response and emerging management needs.

6.13. Integrated River Basin Management Framework

The four assessment frameworks collectively support all major components of Integrated River Basin Management (IRBM).

Table 6.2: Contribution of Assessment Components and Management Stages

Management Activity	River Classification	PRGI	RHA	FACLWS	PMLWS
River inventory	✓				
Conservation planning	✓	✓			✓
Environmental flows	✓	✓	✓		✓
River restoration	✓		✓	✓	✓
Wetland restoration			✓	✓	✓
Urban river rejuvenation			✓	✓	✓
Basin zoning	✓	✓	✓	✓	✓
Climate adaptation	✓	✓	✓	✓	✓
Long-term monitoring			✓	✓	✓

Note: ✓ indicates that the component contributes to the specified management activity.

Important distinction: PMLWS is not a fifth assessment framework. It is the planning and management stage that uses assessment findings to contribute to Water Security, River Health and Climate Resilience.

6.14. Integration with GRBMP 2.0

The framework aligns closely with the objectives of the Ganga River Basin Management Plan (GRBMP 2.0) by supporting:

- ecological river rejuvenation;
- conservation of high-value river reaches;
- environmental flow implementation;
- river-sensitive land-use planning;
- decentralized wastewater management;
- restoration of tributaries and wetlands;
- basin-scale resilience;
- adaptive governance; and
- evidence-based decision-making.

Rather than replacing existing planning processes, the framework strengthens them by providing standardized scientific assessment tools applicable across all river basins.

6.15. Concluding Remarks

The value of an assessment framework lies not in the production of scores or classifications but in its ability to guide better management decisions. The integrated framework developed in this guideline bridges the gap between scientific understanding and practical river basin management by linking river identity, ecological significance, river health, and the performance of local water systems within a unified decision-support system. Through this integration, river managers can move beyond fragmented sectoral approaches toward holistic, process-based management that balances conservation, restoration, sustainable use, and climate resilience. The framework provides a common foundation for planning, implementation, monitoring, and adaptive management, ensuring that interventions are tailored to the natural characteristics and functional needs of each river system.

7. Longitudinal River Classification and Reach-Based Assessment

7.1. Introduction

Rivers are dynamic systems that undergo continuous changes in their physical, hydrological, geomorphological, ecological, and biogeochemical characteristics from their source to their confluence or estuary. Consequently, no river should be regarded as a homogeneous entity. Rather, each river comprises a sequence of interconnected reaches, each possessing distinct characteristics, processes, ecological functions, and management requirements.

Historically, many river assessments have treated entire rivers as single management units, often masking important spatial variations. Such an approach may overlook pristine headwaters, highly degraded urban stretches, ecologically important tributary junctions, or functionally significant floodplain reaches. Effective river basin management therefore requires assessments at an appropriate spatial scale that reflects natural variability.

The integrated framework presented in this guideline adopts the **river reach** as the fundamental unit of assessment. Each reach is characterized using the River Classification Framework (Chapter 2), evaluated for intrinsic ecological significance through the **Pristine River Grading Index (PRGI)** (Chapter 3), assessed for present condition using the **River Health Assessment (RHA)** (Chapter 4), and supported by the **Functional Assessment and Classification of Local Water Systems** (Chapter 5). This chapter describes the methodology for delineating river reaches and integrating these assessments into a longitudinal river profile for basin-scale planning.

7.2. Rationale for Reach-Based Assessment

River characteristics change continuously along their course due to variations in geology, topography, climate, tributary inflows, groundwater interaction, land use, and human interventions. Assessing the river as a single unit can obscure these transitions and lead to generalized management recommendations. **Reach-based assessment offers several advantages:**

- Captures spatial variability in river processes.
- Identifies localized degradation and ecological hotspots.
- Enables targeted conservation and restoration.
- Supports environmental flow assessment.
- Facilitates monitoring of changes over time.
- Aligns with River Styles and fluvial geomorphology principles.

Apply RCF, PRGI and RHA to the main-river reach. Apply FACLWS to individual Local Water Systems and their connected networks, and link each diagnosis to the relevant receiving reach. These linked spatial units provide the basis for coordinated Reach-Specific River Management and PMLWS.

7.3. Concept of River Reaches

A **river reach** is a relatively homogeneous Section of a river within which hydrological, geomorphological, ecological, and functional characteristics remain broadly similar. **Reach boundaries are established where significant changes occur in one or more of the following:**

- Valley setting
- Channel morphology
- River slope
- Geology
- Major tributary confluences
- Flow regime
- Floodplain characteristics
- Land use
- Urbanization
- Hydraulic structures (dams, barrages, weirs)
- Ecological characteristics

Reach lengths may vary from a few kilometres in mountainous terrain to several tens of kilometres in alluvial plains.

7.4. Hierarchical Spatial Framework

For basin planning, river systems should be assessed using a hierarchical spatial framework.

Table 7.1: Hierarchy of Assessment Units

Level	Assessment Unit	Typical Scale	Primary Purpose
I	River Basin	Entire basin	Strategic planning
II	Sub-basin	Major tributary basin	Basin management
III	River	Main river	Longitudinal assessment
IV	Reach	Homogeneous river segment	Functional assessment
V	Site	Local investigation	Project planning

This hierarchy enables assessment results to be aggregated from individual reaches to the basin scale.

7.5. Criteria for Reach Delineation

Reach boundaries should be based on natural transitions rather than administrative boundaries. **The principal delineation criteria include:**

Hydrological Criteria

- Major tributary confluences
- Significant changes in discharge
- Transition from perennial to seasonal flow
- Major groundwater inputs
- Environmental flow change

Geomorphological Criteria

- Change in valley confinement
- Channel pattern transition
- Bed material changes
- Slope breaks
- Floodplain development

Ecological Criteria

- Major habitat transitions
- Riparian vegetation changes
- Wetland associations
- Biodiversity hotspots

Anthropogenic Criteria

- Dams
- Barrages
- Urban areas
- Industrial clusters
- Major abstractions
- Channelization

Where multiple criteria coincide, a single reach boundary should be established.

7.6. Reach Characterization

Each reach should be described using a standardized format.

Table 7.2: Standard Reach Description

Attribute	Description
Reach ID	Unique identification code
River Name	Name of river
Reach Name	Local name
Start Chainage	River kilometre
End Chainage	River kilometre
Reach Length	km
Elevation Range	m
River Classification Code	As per Chapter 2
River Style	Geomorphic description
Dominant Land Use	Forest, agriculture, urban, etc.
Major Tributaries	Important inflows
Key Pressures	Major stressors

7.7. Integrated Reach Assessment

The assessment and management stages for a river reach and its associated Local Water Systems are summarized below.

Table 7.3: Reach Assessment Framework

Component / Management Stage	Output
River Classification	River type and classification code
PRGI	Intrinsic Ecological Significance Tier
RHA	River Health Class and dominant deviations
FACLWS	Functional condition and deficits of associated LWS
Integrated Diagnosis	combined interpretation of pressures, causes, functional deficits and management requirements
Reach-Specific River Management	Reach objectives and management plan
PMLWS	Local Water-System network plan
Implementation and Monitoring	Implemented measures and performance evidence
Reassessment and Adaptive Management	Updated diagnosis and revised management priorities

This integrated assessment provides a complete characterization of every river reach.

7.8. Longitudinal River Profile

The reach assessments are combined to develop a **Longitudinal River Profile**, illustrating changes in river characteristics and management priorities from source to mouth.




Figure 7.1: Typical Longitudinal Profile of a River

This profile provides an immediate visualization of changing ecological significance, functional health, and management needs.

7.9. Reach Prioritization

Assessment results enable reaches to be prioritized according to management objectives.

Table 7.4: Reach Priority Categories

Priority	Typical Characteristics	Management Focus
Priority I	High PRGI, High RHA	Conservation
Priority II	High PRGI, Low RHA	Ecological restoration
Priority III	Moderate PRGI, Moderate RHA	Functional enhancement
Priority IV	Low PRGI, Low RHA	Rehabilitation and pollution control

This prioritization facilitates efficient allocation of financial and institutional resources.

7.10. Reach Report Card

For communication with planners and stakeholders, each reach should be summarized in a standardized **River Reach Report Card**.

Table 7.5: River Reach Report Card

Item	Assessment
River Classification	
River Classification Code	
River Style	
PRGI Grade	
RHA Class	
LWS Category	
Major Stressors	
Restoration Priority	
Recommended Actions	

The report card provides a concise summary suitable for basin management plans, annual monitoring reports, and decision-support systems.

7.11. GIS-Based River Assessment

The integrated framework is designed for implementation within a Geographic Information System (GIS). Each river reach should be represented as a geospatial feature linked to an attribute database containing:

- River Classification Code
- River Style
- PRGI Grade
- RHA Class
- LWS Categories
- Major stressors
- Monitoring stations
- Restoration projects
- Environmental flow requirements
- Land use
- Protected areas

GIS integration enables spatial analysis, visualization, monitoring, and decision support at basin, sub-basin, and reach scales.

7.12. Applications

Longitudinal and reach-based assessment supports:

- Basin planning
- River restoration
- Environmental flow assessment
- Floodplain management
- Climate adaptation
- Biodiversity conservation
- Monitoring of river health trends
- Infrastructure planning
- Prioritization of investments
- Preparation of District River Management Plans (DRMPs)

It also facilitates communication among planners, scientists, administrators, and local communities.

7.13. Integration with GRBMP 2.0

Reach-based assessment provides the operational framework for implementing **GRBMP 2.0**. It enables interventions to be tailored to the specific characteristics and needs of individual river reaches rather than applying uniform measures across an entire river.

This approach supports:

- Reach-specific restoration plans.
- Environmental flow allocation by reach.
- Tributary-specific management.
- Identification of ecological hotspots.
- Integration with district and basin-level planning.
- Performance monitoring through repeat assessments.

7.14. Concluding Remarks

The river reach is the most appropriate operational unit for integrated river assessment because it captures the natural variability of river systems while remaining practical for planning and management. The river reach is the operational unit for River Classification, PRGI and RHA. Associated Local Water Systems and their connected networks constitute the operational units for FACLWS and PMLWS, with their diagnosis and management linked to the relevant receiving river reach. Managers can develop a comprehensive longitudinal understanding of river systems and prioritize interventions based on scientific evidence reach by reach.

This chapter provides the bridge between the conceptual assessment frameworks and their practical application. It lays the foundation for standardized river inventories, GIS-enabled decision support, and adaptive management across river basins.

8. Assessment Methodology and Implementation Protocols

8.1. Introduction

The scientific value of any river assessment framework depends not only on the robustness of its conceptual foundation but also on the consistency and transparency of its implementation. Uniform assessment procedures are essential to ensure that results generated by different organizations, investigators, and river basin authorities remain comparable across river basins and over time.

The four frameworks presented in this guideline namely, River Classification, Pristine River Grading Index (PRGI), River Health Assessment (RHA), and Functional Assessment and Classification of Local Water Systems (FACLWS) are intended for nationwide application. This chapter provides standardized methodologies for data collection, field investigations, indicator selection, scoring, quality assurance, reporting, and periodic reassessment, thereby ensuring scientific consistency and reproducibility.

8.2. Implementation Framework

Implementation follows a structured sequence comprising eight stages.




Figure 8.1 Workflow for River Basin Planning

Each stage builds upon the previous one and should be documented to maintain transparency.

8.3. Stage I – Preliminary Desk Study

The assessment begins with a comprehensive review of available information. The desk study should compile:

- Topographic maps
- River basin maps
- Satellite imagery
- Geological maps
- Land use and land cover maps
- Hydrological records
- Water quality data
- Biodiversity inventories
- Wetland inventories
- Existing environmental flow studies
- Previous river health assessments
- District and basin management plans

The desk study establishes the preliminary understanding of river characteristics and identifies data gaps requiring field verification.

8.4. Stage II – River Segmentation

Using the criteria described in Chapter 7, the river should be divided into homogeneous reaches. Reach delineation should consider:

- Valley setting
- Geology
- Hydrology
- Channel morphology
- Tributary confluences
- Major hydraulic structures
- Land-use transitions
- Urban influence
- Floodplain characteristics

Each reach shall receive a unique identification code to facilitate GIS integration and long-term monitoring.

8.5. Stage III – Field Investigation

Field investigations provide the primary data required for assessment. The objectives are to:

- verify desk study findings;
- characterize river reaches;
- assess physical and ecological conditions;

- document anthropogenic pressures; and
- identify restoration opportunities.

Field surveys should include:

Hydrological Observations

- Flow characteristics
- Water abstractions
- Groundwater interactions
- Floodplain connectivity

Geomorphological Survey

- Channel pattern
- Bed material
- Bank stability
- Floodplain condition
- River Styles verification

Water Quality Assessment

- In-situ measurements (temperature, pH, EC, dissolved oxygen)
- Sampling for laboratory analysis
- Visual observations of pollution

Ecological Survey

- Riparian vegetation
- Aquatic habitats
- Fish and macroinvertebrates
- Wetlands
- Invasive species

Socio-economic Observations

- Water uses
- Religious and cultural sites
- Livelihood dependence
- Community interactions

Photographic documentation and GPS coordinates should accompany all observations.

8.6. Stage IV – Data Compilation and Validation

Data from field surveys, monitoring stations, laboratory analyses, and secondary sources should be integrated into a standardized database. Each dataset should include:

- Source
- Date
- Spatial reference
- Methodology

- Reliability
- Limitations

Where conflicting information exists, preference should be given to the most recent, scientifically reliable, and spatially representative data.

8.7. Stage V – Assessment Procedures

Apply the four complementary assessment components below, using RCF identity and reference context to interpret PRGI and RHA, and linking FACLWS diagnoses to the appropriate receiving reaches.

Step 1: River Classification

Determine:

- Hydrological class
- Geomorphological class
- Sediment regime
- Hydrochemical class
- Source category

Assign the integrated River Classification Code.

Step 2: PRGI

Evaluate:

- Hydrology
- Biodiversity
- Uniqueness
- Ecosystem services

Calculate the composite PRGI score and assign the ecological grade.

Step 3: RHA

Assess H, G, S, W, B and C; compute the weighted RHA Score and assign RHA Class A–E.

Step 4: FACLWS

Assess associated LWS identity, expected functions, present functional condition, deficits and river influence.

Step 5: Subsequent diagnosis and management

Integrate RCF, PRGI, RHA and FACLWS outputs through Integrated Diagnosis. Use this diagnosis to prepare distinct, coordinated Reach-Specific River Management and PMLWS plans. Implement and monitor the plans, then reassess and adapt both diagnosis and management.

8.8. Scoring Protocol

Scoring and classification are component-specific. RCF assigns descriptive classes and a five-element code; it does not assign a health score. PRGI uses the intrinsic-significance scale 5 Exceptional, 4 High, 3 Moderate, 2 Low, 1 Very Low, with the formula, weights and Tier

thresholds in Chapter 3. RHA uses the reference-condition scale and six-component weighting in Chapter 4. FACLWS reports a functional diagnosis against the natural or intended role of the Local Water System; any numerical aggregation requires the declared system-specific scoring schedule specified in Section 5.10. Present degradation shall not reduce a PRGI score or alter intrinsic RCF identity; anthropogenic regulation is recorded separately.

8.9. Data Quality Assurance

Reliable assessments require systematic quality assurance. The following principles should be adopted:

Standardized Methods: All measurements should follow nationally accepted protocols.

Field Verification: Remote sensing interpretations should be validated through field observations.

Replicability: Assessment procedures should be repeatable by independent investigators.

Documentation: All assumptions, expert judgments, and data limitations should be clearly documented.

Independent Review: Major basin assessments should undergo peer review before publication.

8.10. GIS-Based Assessment System

All assessment results should be incorporated into a GIS platform. The spatial database should include:

- River reaches
- River Classification Codes
- PRGI Tiers
- RHA classes
- FACLWS functional condition/deficits
- PMLWS management responses/plans
- Monitoring stations
- Wetlands
- Tributaries
- Hydraulic structures
- Pollution sources
- Restoration projects

GIS integration enables basin-wide visualization, spatial analysis, and monitoring.

8.11. River Reach Report Card

Each assessed reach should be summarized in a standardized report card.

Table 8.2: River Reach Report Card

Item	Description
River Name	
Reach ID	
River Classification Code	
River Style	
PRGI Tier	
RHA Class	
Dominant RHA Deviations	
Associated LWS	
FACLWS Functional Diagnosis	
PMLWS Response	
Main River Management Response	
Major Pressures	
Restoration Priority	
Recommended Actions	
Monitoring Frequency	

This report card forms the basic management unit for river basin planning.

8.12. Basin Summary Dashboard

For basin-level reporting, the results from all reaches should be consolidated into a dashboard.

Table 8.3: Basin Assessment Dashboard

Assessment Component	Basin Summary
Total River Length	
Number of Reaches	
PRGI Tier Distribution	
RHA Class Distribution	
LWS Functional-Deficit Summary	
PMLWS Intervention/Management Priorities	
Priority Restoration Reaches	
Conservation Reaches	
Major Stressors	
Recommended Investments	

The dashboard provides decision-makers with an overview of basin status and priorities.

8.13. Monitoring Frequency

Monitoring intervals should reflect the sensitivity of the river system and the intensity of management interventions.

Table 8.4: Recommended Monitoring Frequency

Assessment Component	Frequency
River Classification	Every 10 years or after major geomorphic changes
PRGI	Every 10 years or following significant ecological changes
RHA	Every 3–5 years
FACLWS	Every 2–3 years
Water Quality	As per national monitoring programme
Hydrology	Continuous where feasible
Biodiversity	Every 3–5 years

Periodic reassessment allows managers to evaluate trends and the effectiveness of interventions.

8.14. Capacity Building

Successful implementation requires trained multidisciplinary teams comprising:

- Hydrologists
- Geomorphologists
- Ecologists
- Environmental engineers
- Water quality specialists
- GIS analysts
- Social scientists
- River basin planners

Training should cover:

- assessment methodologies;
- field survey techniques;
- scoring procedures;
- GIS applications;
- quality assurance; and
- reporting standards.

8.15. Reporting Standards

Assessment reports should include:

- Executive Summary
- Basin Description
- River Segmentation
- River Classification
- PRGI Results
- RHA Results
- FACLWS Results

- Integrated Diagnosis
- Reach-Specific River Management Plan
- PMLWS Plan
- GIS Maps and Reach Report Cards
- Implementation and Monitoring Plan
- Periodic Reassessment and Adaptive-Management Requirements
- Data Sources, Uncertainty and Limitations

Standardized reporting ensures comparability across river basins.

8.16. Digital Decision Support System

The assessment framework is well suited for implementation through a web-based **River Basin Decision Support System (DSS)**.

Core modules may include:

- GIS Viewer
- Reach Database
- Assessment Calculator
- PRGI Module
- RHA Module
- FACLWS Module
- Integrated Diagnosis Module
- Reach-Specific River Management Module
- PMLWS Decision-Support and Planning Module
- Implementation Tracking Module
- Monitoring and Reassessment Dashboard
- Adaptive-Management Decision-Support Module

Such a DSS would facilitate data sharing, periodic updates, and evidence-based decision-making.

8.17. Concluding Remarks

The methodologies presented in this chapter provide a standardized operational framework for implementing the integrated river assessment system. By combining rigorous field investigations, consistent scoring protocols, GIS-based data management, quality assurance, and periodic reassessment, the framework ensures that River Classification, PRGI, RHA, and FACLWS can be applied uniformly across diverse river basins.

The resulting assessments are synthesised through Integrated Diagnosis to guide Reach-Specific River Management and PMLWS. Implementation, monitoring and periodic reassessment provide the evidence required for adaptive revision of management measures and priorities.

9. Implementation Framework for Integrated River Basin Management

Towards a National Framework for River Assessment, Conservation, Restoration and Sustainable Management

"A river cannot be restored merely by improving its water quality, nor can it be conserved solely by protecting isolated reaches. Sustainable river management requires an integrated understanding of river identity, ecological significance, functional health, and the contribution of supporting aquatic systems. The Integrated River Assessment Framework provides the scientific foundation for achieving this objective."

9.1. Introduction

9.1.1. From Assessment to Action

The preceding chapters have established the scientific principles, conceptual foundations, and operational methodologies of the Integrated River Assessment Framework (IRAF). Together, the River Classification Framework (RCF), the Pristine River Grading Index (PRGI), the River Health Assessment (RHA) and FACLWS constitute the assessment and diagnostic architecture of IRAF. PMLWS is the subsequent planning and management stage for associated Local Water Systems and is not a fifth assessment framework.

However, scientific assessment alone cannot restore rivers. Assessments become meaningful only when they are translated into planning, policy, investment, implementation, monitoring, and adaptive management. River restoration is therefore not an end product of assessment but the outcome of an iterative process that integrates scientific understanding with governance, institutional coordination, engineering interventions, ecological restoration, stakeholder participation, and long-term monitoring.

Historically, river management in many parts of the world has evolved through sectoral approaches focusing separately on water resources development, flood management, pollution control, irrigation, hydropower, navigation, biodiversity conservation, or urban infrastructure. While each of these sectors addresses important aspects of river systems, they often operate independently, resulting in fragmented planning, duplication of efforts, conflicting objectives, and inefficient resource utilization. Such fragmented approaches rarely achieve sustainable river rejuvenation because they fail to recognize rivers as integrated socio-ecological systems.

The Integrated River Assessment Framework has been developed specifically to overcome these limitations. Rather than treating rivers solely as hydraulic systems, water resources,

drainage channels, or ecological habitats, the framework recognizes rivers as dynamic, interconnected systems whose hydrological, geomorphological, ecological, environmental, social, cultural, and economic functions are inseparable.

Accordingly, implementation of the framework requires a corresponding transformation in river governance—from project-based interventions to basin-scale planning, from isolated engineering works to integrated ecosystem management, from reactive pollution control to preventive river stewardship, and from static planning to adaptive management.

This chapter provides the strategic framework for translating scientific assessments into actionable river basin management programmes. It establishes the institutional, planning, technological, and governance mechanisms necessary for implementing the Integrated River Assessment Framework across river basins of varying scales and complexities.

9.1.2. Need for an Integrated Implementation Framework

River basin management has entered a new era characterized by increasing environmental uncertainty, rapid urbanization, changing climate, competing water demands, and growing societal expectations. Rivers are now expected not only to supply water but also to sustain biodiversity, regulate floods, recharge groundwater, support livelihoods, preserve cultural heritage, provide recreational opportunities, maintain ecosystem services, and contribute to climate resilience.

These expanding expectations cannot be met through conventional river management approaches that primarily focus on individual sectors or isolated projects. Instead, they require integrated planning supported by comprehensive scientific assessment.

The Integrated River Assessment Framework fulfils this need by establishing a common scientific language for river assessment and management. River Classification defines the physical identity of river systems; PRGI identifies reaches of high intrinsic ecological significance requiring long-term conservation; RHA evaluates present functional performance and identifies restoration needs; while FACLWS assesses the contribution of associated aquatic systems that collectively sustain basin resilience.

Together, these frameworks provide a robust scientific basis for evidence-based planning, enabling managers to distinguish between conservation, protection, restoration, rehabilitation, and sustainable development priorities. More importantly, they provide a common platform through which scientists, engineers, administrators, planners, policymakers, and local communities can communicate using consistent terminology and standardized assessment methodologies.

9.1.3. Purpose of this Chapter

Unlike the preceding chapters, which focus on scientific concepts and assessment methodologies, this chapter focuses on implementation. Its purpose is to demonstrate how the assessment framework can be embedded within the broader process of Integrated River Basin Management (IRBM), thereby transforming scientific information into practical decision support.

Specifically, this chapter aims to:

- establish a strategic roadmap for implementing the Integrated River Assessment Framework at national, basin, sub-basin, river, reach, and local water system scales;
- provide guidance on institutional arrangements and governance mechanisms necessary for coordinated implementation;
- integrate scientific assessments with planning, investment, restoration, monitoring, and adaptive management;
- promote standardized implementation across river basins to ensure consistency and comparability of assessments;
- identify opportunities for digital transformation through GIS, remote sensing, artificial intelligence, and decision-support systems; and
- articulate a long-term national vision for river assessment, conservation, restoration, and sustainable management.



Figure 9.1: Evolution of River Basin Management

In doing so, the chapter serves as a bridge between scientific assessment and practical river basin management, thereby completing the conceptual framework presented in this volume.

9.2. From Assessment to Integrated River Basin Management

9.2.1. Assessment as the Foundation of River Management

Effective river management begins with understanding the river itself. Without a sound understanding of river identity, ecological significance, functional condition, and associated aquatic systems, management interventions are likely to be fragmented, inefficient, or even counterproductive.

The Integrated River Assessment Framework therefore adopts the principle of "**Assessment before Intervention.**" This principle recognizes that every river possesses unique physical characteristics, ecological attributes, and functional processes that should guide management decisions. Restoration measures appropriate for a Himalayan River may be unsuitable for an alluvial river; similarly, interventions designed for a pristine headwater stream may be inappropriate for a highly urbanized river reach. Standardized assessment provides the scientific basis for identifying these differences and tailoring interventions accordingly.

Assessment is thus not merely a diagnostic exercise but the foundation upon which planning, investment, implementation, monitoring, and adaptive management are built.

9.2.2. Linking Assessment with the River Basin Planning Cycle

The Integrated River Assessment Framework is designed to support every stage of the river basin planning cycle.

Initially, River Classification establishes the natural identity of the river and defines the reference conditions against which future changes are evaluated. PRGI identifies reaches possessing high ecological significance that require conservation. RHA evaluates current functional performance and identifies restoration priorities, while FACLWS extends the assessment to floodplain wetlands, tributaries, springs, ponds, paleochannels, and engineered aquatic systems.

The outputs of these assessments are subsequently synthesized to identify management priorities, formulate restoration strategies, allocate investments, design monitoring programmes, and evaluate implementation outcomes. Periodic reassessment allows managers to determine whether interventions are producing measurable improvements in river condition and, where necessary, modify management strategies accordingly.

The planning cycle therefore becomes an iterative process characterized by continuous learning and adaptive management rather than a one-time planning exercise.




Figure 9.2: Integrated River Assessment-to-Management Pathway—From River Classification and Functional Diagnosis to Implementation, Monitoring and Adaptive Management

The framework integrates four complementary assessment components—River Classification (RCF), Pristine River Grading Index (PRGI), River Health Assessment (RHA), and Functional Assessment and Classification of Local Water Systems (FACLWS)—into an Integrated Diagnosis. This informs two complementary planning and management pathways: reach-specific river management and Planning and Management of Local Water Systems (PMLWS). Implementation is followed by monitoring, reassessment and adaptive management, creating a continuous feedback cycle for improving River Health, Water Security and Climate Resilience.

9.2.3. Integrated River Basin Management Philosophy

The philosophy underlying the Integrated River Assessment Framework is founded on the recognition that rivers should be managed as living systems rather than as isolated water conveyance channels. Hydrology, geomorphology, sediment transport, water quality, habitats, biodiversity, floodplains, wetlands, groundwater, and human communities together constitute an interconnected socio-ecological system whose components continuously influence one another.

Consequently, river basin management should seek to restore and sustain the functioning of the entire river system rather than addressing individual problems in isolation. Pollution control, environmental flows, floodplain restoration, biodiversity conservation, groundwater recharge, wastewater reuse, and nature-based solutions should therefore be viewed as complementary components of a unified management strategy rather than independent programmes.

This integrated philosophy provides the conceptual foundation for GRBMP 2.0 and aligns river basin planning with contemporary principles of ecosystem-based management, adaptive governance, and sustainable development.

Table 9.1: Transition from Conventional River Management to Integrated River Basin Management

Conventional Approach	Integrated River Assessment Framework
Project-oriented	Basin-oriented
Water quality focused	River system focused
Administrative boundaries	Hydrological boundaries
Sectoral planning	Integrated planning
Reactive pollution control	Preventive ecosystem management
Engineering solutions	Hybrid engineering and nature-based solutions
Periodic project evaluation	Continuous adaptive management
Single-sector decision-making	Multi-disciplinary governance
Static plans	Dynamic, evidence-based planning

9.3. Guiding Principles for Implementation

Successful implementation of the Integrated River Assessment Framework depends upon a set of guiding principles that ensure consistency, scientific integrity, and long-term sustainability across all river basins.

These principles provide the philosophical foundation for river assessment, planning, restoration, monitoring, and adaptive management.

Principle 1 – River Identity Before Intervention

Every river possesses unique hydrological, geomorphological, ecological, and cultural characteristics. Management interventions shall therefore be based on River Classification and reference conditions rather than standardized engineering prescriptions.

Principle 2 – Conservation Before Restoration

River reaches exhibiting high PRGI and good RHA should receive priority for conservation and protection, thereby preventing future degradation and reducing long-term restoration costs.

Principle 3 – Functional Integrity

Management should focus on restoring the natural functioning of rivers rather than merely improving individual environmental indicators such as water quality.

Principle 4 – Basin-scale Planning

Planning shall follow hydrological boundaries at basin, sub-basin, river-reach and connected Local Water-System scales, while implementation shall be organized through appropriate administrative units and institutions.

Principle 5 – Reach-based Management

The river reach shall constitute the principal operational unit for main-river classification, PRGI, RHA and reach-level management. Individual and connected networks of Local Water Systems shall constitute the operational units for FACLWS and PMLWS.

Principle 6 – Integration of Local Water Systems

Local Water Systems shall be assessed through FACLWS and planned and managed through PMLWS as distributed hydrological and ecological infrastructure contributing to Water Security, River Health and Climate Resilience.

Principle 7 – Science-based Decision Making

Planning, investment, and restoration priorities shall be guided by standardized scientific assessments rather than subjective judgement alone.

Principle 8 – Adaptive Management

River management shall be treated as a continuous learning process supported by periodic reassessment, monitoring, and evidence-based revision of management strategies.

Principle 9 – Multi-stakeholder Participation

Implementation should involve government agencies, river basin organizations, local authorities, academic institutions, civil society organizations, industries, and river-dependent communities.

Principle 10 – Climate Resilience and Sustainability

River basin management should enhance resilience to climate variability while ensuring long-term ecological sustainability and water security.

Principle 11 – Digital Transformation

Implementation should progressively adopt GIS, Earth observation, artificial intelligence, sensor networks, and digital decision-support systems to improve assessment efficiency, transparency, and responsiveness.

Principle 12 – Knowledge Integration

River management should integrate scientific research, traditional ecological knowledge, local experience, and emerging technologies to continuously improve understanding and management of river systems.




Figure 9.3: Twelve Guiding Principles of the Integrated River Assessment Framework

Concluding Note

The concepts presented in this Volume establish the philosophical and strategic foundation for implementing the Integrated River Assessment Framework. Volume II demonstrates the operational application of IRAF through the Tier I Ganga River case, including River Classification, PRGI, RHA,

FACLWS and the translation of diagnosis into coordinated Reach-Specific River Management and PMLWS. The framework is also applicable to other river systems, including the Hindon.